



BN

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 471/04, 401/06, 401/14, 519/00, A61K 31/415, 31/44 // (C07D 519/00, 498:00, 471:00) (C07D 519/00, 491:00, 471:00) (C07D 519/00, 471:00, 471:00)</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/31099</b> (43) International Publication Date: <b>24 June 1999 (24.06.99)</b>																																													
(21) International Application Number: <b>PCT/US98/26568</b> (22) International Filing Date: <b>14 December 1998 (14.12.98)</b> (30) Priority Data: <table><tr><td>60/069,910</td><td>17 December 1997 (17.12.97)</td><td>US</td></tr><tr><td>60/079,197</td><td>24 March 1998 (24.03.98)</td><td>US</td></tr><tr><td>60/079,944</td><td>30 March 1998 (30.03.98)</td><td>US</td></tr><tr><td>60/080,397</td><td>2 April 1998 (02.04.98)</td><td>US</td></tr><tr><td>60/083,251</td><td>27 April 1998 (27.04.98)</td><td>US</td></tr><tr><td>9810182.7</td><td>13 May 1998 (13.05.98)</td><td>GB</td></tr><tr><td>9810882.2</td><td>20 May 1998 (20.05.98)</td><td>GB</td></tr><tr><td>9810892.1</td><td>20 May 1998 (20.05.98)</td><td>GB</td></tr><tr><td>9811283.2</td><td>26 May 1998 (26.05.98)</td><td>GB</td></tr><tr><td>9812686.5</td><td>12 June 1998 (12.06.98)</td><td>GB</td></tr><tr><td>60/092,588</td><td>13 July 1998 (13.07.98)</td><td>US</td></tr><tr><td>60/092,624</td><td>13 July 1998 (13.07.98)</td><td>US</td></tr><tr><td>60/099,948</td><td>11 September 1998 (11.09.98)</td><td>US</td></tr><tr><td>9822331.6</td><td>13 October 1998 (13.10.98)</td><td>GB</td></tr><tr><td>9822701.0</td><td>16 October 1998 (16.10.98)</td><td>GB</td></tr></table> (71) Applicant (for all designated States except US): <b>MERCK &amp; CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</b>			60/069,910	17 December 1997 (17.12.97)	US	60/079,197	24 March 1998 (24.03.98)	US	60/079,944	30 March 1998 (30.03.98)	US	60/080,397	2 April 1998 (02.04.98)	US	60/083,251	27 April 1998 (27.04.98)	US	9810182.7	13 May 1998 (13.05.98)	GB	9810882.2	20 May 1998 (20.05.98)	GB	9810892.1	20 May 1998 (20.05.98)	GB	9811283.2	26 May 1998 (26.05.98)	GB	9812686.5	12 June 1998 (12.06.98)	GB	60/092,588	13 July 1998 (13.07.98)	US	60/092,624	13 July 1998 (13.07.98)	US	60/099,948	11 September 1998 (11.09.98)	US	9822331.6	13 October 1998 (13.10.98)	GB	9822701.0	16 October 1998 (16.10.98)	GB	(72) Inventors; and (75) Inventors/Applicants (for US only): <b>DUGGAN, Mark, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MEISSNER, Robert, S. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HUTCHINSON, John, H. [CA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HALCZENKO, Wasył [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ASKEW, Ben, C. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). COLEMAN, Paul, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PATANE, Michael, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WANG, Jiabing [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</b> (74) Common Representative: <b>MERCK &amp; CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</b> (81) Designated States: <b>AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
60/069,910	17 December 1997 (17.12.97)	US																																														
60/079,197	24 March 1998 (24.03.98)	US																																														
60/079,944	30 March 1998 (30.03.98)	US																																														
60/080,397	2 April 1998 (02.04.98)	US																																														
60/083,251	27 April 1998 (27.04.98)	US																																														
9810182.7	13 May 1998 (13.05.98)	GB																																														
9810882.2	20 May 1998 (20.05.98)	GB																																														
9810892.1	20 May 1998 (20.05.98)	GB																																														
9811283.2	26 May 1998 (26.05.98)	GB																																														
9812686.5	12 June 1998 (12.06.98)	GB																																														
60/092,588	13 July 1998 (13.07.98)	US																																														
60/092,624	13 July 1998 (13.07.98)	US																																														
60/099,948	11 September 1998 (11.09.98)	US																																														
9822331.6	13 October 1998 (13.10.98)	GB																																														
9822701.0	16 October 1998 (16.10.98)	GB																																														
(54) Title: <b>INTEGRIN RECEPTOR ANTAGONISTS</b> (57) Abstract <p>The present invention relates to compounds and derivatives thereof, their synthesis, and their use as integrin receptor antagonists. More particularly, the compounds of the present invention are antagonists of the integrin receptors <math>\alpha v \beta 3</math>, <math>\alpha v \beta 5</math> and/or <math>\alpha v \beta 6</math> and are useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, wound healing, viral disease, and tumor growth and metastasis.</p>																																																

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**TITLE OF THE INVENTION**  
**INTEGRIN RECEPTOR ANTAGONISTS**

5

**CROSS-REFERENCE TO RELATED APPLICATIONS**

The present invention is related to U.S. provisional applications Serial No. 60/069,910, filed December 17, 1997; 60/083,251, filed April 27, 1998; 60/092,588, filed July 13, 1998; 60/079,197, filed March 24, 1998; 60/079,944, filed March 30, 1998; 60/080,397, filed April 2, 1998; 60/092,624, filed July 13, 1998; and 60/099,948, filed September 11, 1998; the contents of each of which are hereby incorporated by reference.

10

**FIELD OF THE INVENTION**

15

The present invention relates to compounds and derivatives thereof, their synthesis, and their use as integrin receptor antagonists. More particularly, the compounds of the present invention are antagonists of the integrin receptors  $\alpha v \beta 3$ ,  $\alpha v \beta 5$ , and/or  $\alpha v \beta 6$  and are useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, wound healing, viral disease, tumor growth, and metastasis.

20

**BACKGROUND OF THE INVENTION**

25

It is believed that a wide variety of disease states and conditions can be mediated by acting on integrin receptors and that integrin receptor antagonists represent a useful class of drugs. Integrin receptors are heterodimeric transmembrane proteins through which cells attach and communicate with extracellular matrices and other cells (See S.B. Rodan and G.A. Rodan, "Integrin Function In Osteoclasts", *Journal of Endocrinology*, Vol. 154, S47- S56 (1997), which is incorporated by reference herein in its entirety).

30

In one aspect of the present invention, the compounds herein are useful for inhibiting bone resorption. Bone resorption is mediated by the action of cells known as osteoclasts. Osteoclasts are

35

large multinucleated cells of up to about 400  $\mu$ m in diameter that resorb mineralized tissue, chiefly calcium carbonate and calcium phosphate, in vertebrates. Osteoclasts are actively motile cells that migrate along the surface of bone, and can bind to bone, secrete necessary acids and proteases, thereby causing the actual resorption of mineralized tissue from the bone. More specifically, osteoclasts are believed to exist in at least two physiological states, namely, the secretory state and the migratory or motile state. In the secretory state, osteoclasts are flat, attach to the bone matrix via a tight attachment zone (sealing zone), become highly polarized, form a ruffled border, and secrete lysosomal enzymes and protons to resorb bone. The adhesion of osteoclasts to bone surfaces is an important initial step in bone resorption. In the migratory or motile state, the osteoclasts migrate across bone matrix and do not take part in resorption until they again attach to bone.

Integrins are involved in osteoclast attachment, activation and migration. The most abundant integrin in osteoclasts, e.g., in rat, chicken, mouse and human osteoclasts, is an integrin receptor known as  $\alpha_v\beta_3$ , which is thought to interact in bone with matrix proteins that contain the RGD sequence. Antibodies to  $\alpha_v\beta_3$  block bone resorption *in vitro* indicating that this integrin plays a key role in the resorptive process. There is increasing evidence to suggest that  $\alpha_v\beta_3$  ligands can be used effectively to inhibit osteoclast mediated bone resorption *in vivo* in mammals.

The current major bone diseases of public concern are osteoporosis, hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, periarticular erosions in rheumatoid arthritis, Paget's disease, immobilization-induced osteopenia, and glucocorticoid-induced osteoporosis. All of these conditions are characterized by bone loss, resulting from an imbalance between bone resorption, i.e. breakdown, and bone formation, which continues throughout life at the rate of about 14% per year on the average. However, the rate of bone turnover differs from site to site; for example, it is higher in the trabecular bone of the vertebrae and the alveolar bone in the jaws than in the cortices of the long bones. The potential for bone loss is directly related to turnover and can amount to



over 5% per year in vertebrae immediately following menopause, a condition which leads to increased fracture risk.

In the United States, there are currently about 20 million people with detectable fractures of the vertebrae due to osteoporosis. In addition, there are about 250,000 hip fractures per year attributed to osteoporosis. This clinical situation is associated with a 12% mortality rate within the first two years, while 30% of the patients require nursing home care after the fracture.

Individuals suffering from all the conditions listed above would benefit from treatment with agents which inhibit bone resorption.

Additionally,  $\alpha v \beta 3$  ligands have been found to be useful in treating and/or inhibiting restenosis, i.e. recurrence of stenosis after corrective surgery on the heart valve, atherosclerosis, diabetic retinopathy, macular degeneration, and angiogenesis, i.e. formation of new blood vessels. Moreover, it has been postulated that the growth of tumors depends on an adequate blood supply, which in turn is dependent on the growth of new vessels into the tumor; thus, inhibition of angiogenesis can cause tumor regression in animal models (See Harrison's Principles of Internal Medicine, 12th ed., 1991, which is incorporated by reference herein in its entirety). Therefore,  $\alpha v \beta 3$  antagonists which inhibit angiogenesis can be useful in the treatment of cancer by inhibiting tumor growth (See e.g., Brooks et al., Cell, 79:1157-1164 (1994), which is incorporated by reference herein in its entirety).

Moreover, compounds of this invention can also inhibit neovascularization by acting as antagonists of the integrin receptor,  $\alpha v \beta 5$ . A monoclonal antibody for  $\alpha v \beta 5$  has been shown to inhibit VEGF-induced angiogenesis in rabbit cornea and the chick chorioallantoic membrane model (See M.C. Friedlander, et al., Science 270, 1500-1502, (1995), which is incorporated by reference herein in its entirety). Thus, compounds that antagonize  $\alpha v \beta 5$  are useful for treating and preventing macular degeneration, diabetic retinopathy, tumor growth, and metastasis.

Additionally, compounds of the instant invention can inhibit angiogenesis and inflammation by acting as antagonists of the integrin receptor,  $\alpha v \beta 6$ , which is expressed during the later stages of

wound healing and remains expressed until the wound is closed (See Christofidou-Solomidou, et al., "Expression and Function of Endothelial Cell  $\alpha$ v Integrin Receptors in Wound-Induced Human Angiogenesis in Human Skin/SCID Mice Chimeras, American Journal of Pathology, Vol. 151, No. 4, pp. 975-983 (October 1997), which is incorporated by reference herein in its entirety). It is postulated that  $\alpha$ v $\beta$ 6 plays a role in the remodeling of the vasculature during the later stages of angiogenesis. Also,  $\alpha$ v $\beta$ 6 participates in the modulation of epithelial inflammation and is induced in response to local injury or inflammation (See Xiao-Zhu Huang, et al., "Inactivation of the Integrin  $\beta$ 6 Subunit Gene Reveals a Role of Epithelial Integrins in Regulating Inflammation in the Lungs and Skin," Journal of Cell Biology, Vol. 133, No.4, pp. 921-928 (May 1996), which is incorporated by reference herein in its entirety). Accordingly, compounds that antagonize  $\alpha$ v $\beta$ 6 are useful in treating or preventing cancer by inhibiting tumor growth and metastasis.

In addition, certain compounds of this invention antagonize both the  $\alpha$ v $\beta$ 3 and  $\alpha$ v $\beta$ 5 receptors. These compounds, referred to as "dual  $\alpha$ v $\beta$ 3/ $\alpha$ v $\beta$ 5 antagonists," are useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, tumor growth, and metastasis.

In addition, certain compounds of this invention are useful as mixed  $\alpha$ v $\beta$ 3,  $\alpha$ v $\beta$ 5, and  $\alpha$ v $\beta$ 6 receptor antagonists.

It is therefore an object of the present invention to provide compounds which are useful as integrin receptor antagonists.

It is another object of the present invention to provide compounds which are useful as  $\alpha$ v $\beta$ 3 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as  $\alpha$ v $\beta$ 5 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as  $\alpha$ v $\beta$ 6 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as dual  $\alpha$ v $\beta$ 3/ $\alpha$ v $\beta$ 5 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as mixed  $\alpha v \beta 3$ ,  $\alpha v \beta 5$ , and  $\alpha v \beta 6$  receptor antagonists.

It is another object of the present invention to provide  
5 pharmaceutical compositions comprising integrin receptor antagonists.

It is another object of the present invention to provide methods for making the pharmaceutical compositions of the present invention.

It is another object of the present invention to provide  
10 methods for eliciting an integrin receptor antagonizing effect in a mammal in need thereof by administering the compounds and pharmaceutical compositions of the present invention.

It is another object of the present invention to provide compounds and pharmaceutical compositions useful for inhibiting bone  
15 resorption, restenosis, atherosclerosis, inflammation, viral disease, diabetic retinopathy, macular degeneration, angiogenesis, tumor growth, and metastasis.

It is another object of the present invention to provide compounds and pharmaceutical compositions useful for treating  
20 osteoporosis.




It is another object of the present invention to provide methods for inhibiting bone resorption, restenosis, atherosclerosis, inflammation, viral disease, diabetic retinopathy, macular  
degeneration, angiogenesis, tumor growth, and metastasis.

It is another object of the present invention to provide  
25 methods for treating osteoporosis.



These and other objects will become readily apparent from the detailed description which follows.


### 30 SUMMARY OF THE INVENTION

The present invention relates to compounds having a structural formula selected from the group consisting of




5






, and



;

- 5 wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are substituted only with  $R^{10}$  and  $R^{12}$ ;

X is selected from the group consisting of



- 10 a 5- or 6-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one  $R^1$  substituent and the ring carbon atoms are unsubstituted or substituted with one or two  $R^1$  substituents, and
- 15

- a 9- to 14-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or
- 20

substituted with one  $R^1$  substituent and the ring carbon atoms are unsubstituted or substituted with one or two  $R^1$  substituents;

Y is selected from the group consisting of

- 5
- (CH<sub>2</sub>)<sub>m</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>-,
  - 10 -(CH<sub>2</sub>)<sub>m</sub>-SO-(CH<sub>2</sub>)<sub>n</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>n</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-,
  - 15 -(CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-S-(CH<sub>2</sub>)<sub>p</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>-S-(CH<sub>2</sub>)<sub>p</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-S-(CH<sub>2</sub>)<sub>p</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,
  - (CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,
  - 20 -(CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-, and
  - (CH<sub>2</sub>)<sub>m</sub>-Z-(CH<sub>2</sub>)<sub>n</sub>-,

wherein Z is a 3- to 10-membered monocyclic or polycyclic aromatic or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one  $R^1$  substituent and the ring carbon atoms are unsubstituted or substituted with one or two  $R^1$  substituents, and wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than in  $R^4$ , can be substituted by one or two  $R^3$  substituents; and

30 wherein  $R^1$  and  $R^2$  are each independently selected from the group consisting of

hydrogen, halogen, C<sub>1</sub>-10 alkyl, C<sub>3</sub>-8 cycloalkyl,  
C<sub>3</sub>-8 cycloheteroalkyl, C<sub>3</sub>-8 cycloalkyl C<sub>1</sub>-6 alkyl,

C<sub>3-8</sub> cycloheteroalkyl C<sub>1-6</sub> alkyl, aryl, aryl C<sub>1-8</sub> alkyl, amino,  
 amino C<sub>1-8</sub> alkyl, C<sub>1-3</sub> acylamino, C<sub>1-3</sub> acylamino C<sub>1-8</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino, (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-8</sub> alkyl,  
 C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkoxy C<sub>1-6</sub> alkyl, hydroxycarbonyl,  
 5 hydroxycarbonyl C<sub>1-6</sub> alkyl, C<sub>1-3</sub> alkoxycarbonyl,  
 C<sub>1-3</sub> alkoxycarbonyl C<sub>1-6</sub> alkyl, hydroxycarbonyl-  
 C<sub>1-6</sub> alkyloxy, hydroxy, hydroxy C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkyloxy-  
 C<sub>1-6</sub> alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,  
 trifluoroethoxy, C<sub>1-8</sub> alkyl-S(O)<sub>p</sub>, (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 10 C<sub>1-8</sub> alkyloxycarbonylamino, (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>amino, (aryl)<sub>p</sub>amino, aryl C<sub>1-8</sub>  
 alkylsulfonylamino, and C<sub>1-8</sub> alkylsulfonylamino;  
 or two R<sup>1</sup>-substituents, when on the same carbon atom, are taken  
 together with the carbon atom to which they are attached to  
 15 form a carbonyl group;

each R<sup>3</sup> is independently selected from the group consisting of  
 hydrogen,  
 aryl,  
 20 C<sub>1-10</sub> alkyl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 25 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 halogen,  
 hydroxyl,  
 oxo,  
 30 trifluoromethyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,

- C<sub>1-6</sub> alkylcarbonyloxy,  
 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 amino C<sub>1-6</sub> alkyl,  
 5 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 hydroxycarbonyl,  
 10 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 15 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 20 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 25 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 30 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 C<sub>1-8</sub> alkylsulfonylamino,



- arylsulfonylamino,  
C1-8 alkylsulfonylamino C1-6 alkyl,  
arylsulfonylamino C1-6 alkyl,  
aryl C1-6 alkylsulfonylamino,  
5 aryl C1-6 alkylsulfonylamino C1-6 alkyl,  
C1-8 alkoxycarbonylamino,  
C1-8 alkoxycarbonylamino C1-8 alkyl,  
aryloxycarbonylamino C1-8 alkyl,  
aryl C1-8 alkoxycarbonylamino,  
10 aryl C1-8 alkoxycarbonylamino C1-8 alkyl,  
C1-8 alkylcarbonylamino,  
C1-8 alkylcarbonylamino C1-6 alkyl,  
arylcarbonylamino C1-6 alkyl,  
aryl C1-6 alkylcarbonylamino,  
15 aryl C1-6 alkylcarbonylamino C1-6 alkyl,  
aminocarbonylamino C1-6 alkyl,  
(C1-8 alkyl)paminocarbonylamino,  
(C1-8 alkyl)paminocarbonylamino C1-6 alkyl,  
(aryl)paminocarbonylamino C1-6 alkyl,  
20 (aryl C1-8 alkyl)paminocarbonylamino,  
(aryl C1-8 alkyl)paminocarbonylamino C1-6 alkyl,  
aminosulfonylamino C1-6 alkyl,  
(C1-8 alkyl)paminosulfonylamino,  
(C1-8 alkyl)paminosulfonylamino C1-6 alkyl,  
25 (aryl)paminosulfonylamino C1-6 alkyl,  
(aryl C1-8 alkyl)paminosulfonylamino,  
(aryl C1-8 alkyl)paminosulfonylamino C1-6 alkyl,  
C1-6 alkylsulfonyl,  
C1-6 alkylsulfonyl C1-6 alkyl,  
30 arylsulfonyl C1-6 alkyl,  
aryl C1-6 alkylsulfonyl,  
aryl C1-6 alkylsulfonyl C1-6 alkyl,  
C1-6 alkylcarbonyl,  
C1-6 alkylcarbonyl C1-6 alkyl,  
35 arylcarbonyl C1-6 alkyl,

- aryl C<sub>1-6</sub> alkylcarbonyl,  
aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
5 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
10 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl;  
or two R<sup>3</sup> substituents, when on the same carbon atom, are taken  
together with the carbon atom to which they are attached to  
form a carbonyl group or a cyclopropyl group,  
15 wherein any of the alkyl groups of R<sup>3</sup> are either unsubstituted or  
substituted with one to three R<sup>1</sup> substituents,  
and provided that each R<sup>3</sup> is selected such that in the resultant  
compound the carbon atom or atoms to which R<sup>3</sup> is attached is itself  
attached to no more than one heteroatom;  
20 each R<sup>4</sup> is independently selected from the group consisting of  
hydrogen,  
aryl,  
aminocarbonyl,  
25 C<sub>3-8</sub> cycloalkyl,  
amino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl,  
(aryl C<sub>1-5</sub> alkyl)<sub>p</sub>aminocarbonyl,  
hydroxycarbonyl C<sub>1-6</sub> alkyl,  
30 C<sub>1-8</sub> alkyl,  
aryl C<sub>1-6</sub> alkyl,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
(aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
C<sub>1-8</sub> alkylsulfonyl,

- C<sub>1-8</sub> alkoxy carbonyl,  
 aryloxy carbonyl,  
 aryl C<sub>1-8</sub> alkoxy carbonyl,  
 C<sub>1-8</sub> alkyl carbonyl,  
 5 aryl carbonyl,  
 aryl C<sub>1-6</sub> alkyl carbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub> aminocarbonyl,  
 aminosulfonyl,  
 C<sub>1-8</sub> alkyl aminosulfonyl,  
 10 (aryl)<sub>p</sub> aminosulfonyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub> aminosulfonyl,  
 aryl sulfonyl,  
 aryl C<sub>1-6</sub> alkyl sulfonyl,  
 C<sub>1-6</sub> alkyl thiocarbonyl,  
 15 aryl thiocarbonyl, and  
 aryl C<sub>1-6</sub> alkyl thiocarbonyl,

wherein any of the alkyl groups of R<sup>4</sup> are either unsubstituted or substituted with one to three R<sup>1</sup> substituents;

- 20 R<sup>5</sup> and R<sup>6</sup> are each independently selected from the group consisting of  
 hydrogen,  
 C<sub>1-10</sub> alkyl,  
 aryl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 25 aryl-(CH<sub>2</sub>)<sub>r</sub>-S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 30 halogen,  
 hydroxyl,  
 C<sub>1-8</sub> alkyl carbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxy carbonyl,

- (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 5 amino C<sub>1-6</sub> alkyl,  
 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 10 hydroxycarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 15 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 20 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 25 aryl C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 30 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,

- C<sub>1-8</sub> alkylsulfonylamino,  
 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 5 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkoxycarbonylamino,  
 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
 10 aryl C<sub>1-8</sub> alkoxycarbonylamino,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 arylcarbonylamino C<sub>1-6</sub> alkyl,  
 15 aryl C<sub>1-6</sub> alkylcarbonylamino,  
 aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 20 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 25 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylsulfonyl,  
 30 C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 arylsulfonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylcarbonyl,

C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 arylcarbonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 5 C<sub>1-6</sub> alkylthiocarbonylamino,  
 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 10 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl;  
 or R<sup>5</sup> and R<sup>6</sup> are taken together with the carbon atom to which they are  
 15 attached to form a carbonyl group,  
 wherein any of the alkyl groups of R<sup>5</sup> or R<sup>6</sup> are either unsubstituted or  
 substituted with one to three R<sup>1</sup> substituents,  
 and provided that each R<sup>5</sup> and R<sup>6</sup> are selected such that in the resultant  
 compound the carbon atom to which R<sup>5</sup> and R<sup>6</sup> are attached is itself  
 20 attached to no more than one heteroatom;

R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of  
 hydrogen,  
 C<sub>1-10</sub> alkyl,  
 25 aryl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 30 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 halogen,  
 hydroxyl,  
 C<sub>1-8</sub> alkylcarbonylamino,

- aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 5 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 amino C<sub>1-6</sub> alkyl,  
 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 10 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 hydroxycarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 15 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 20 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 25 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 30 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,

- aryl C<sub>1-6</sub> alkylcarbonyloxy,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
C<sub>1-8</sub> alkylsulfonylamino,  
arylcarbonylamino,  
5 arylsulfonylamino,  
C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
arylsulfonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonylamino,  
aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
10 C<sub>1-8</sub> alkoxycarbonylamino,  
C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
aryl C<sub>1-8</sub> alkoxycarbonylamino,  
aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
15 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonylamino,  
aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
20 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
arylaminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
25 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
30 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylsulfonyl,  
C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
arylsulfonyl C<sub>1-6</sub> alkyl,



aryl C<sub>1-6</sub> alkylsulfonyl,  
aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylcarbonyl,  
C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
5 arylcarbonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
10 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
15 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl, and  
C<sub>7-20</sub> polycyclyl C<sub>0-8</sub> alkylsulfonylamino;

wherein any of the alkyl groups of R<sup>7</sup> and R<sup>8</sup> are either unsubstituted or substituted with one to three R<sup>1</sup> substituents,  
20 and provided that each R<sup>7</sup> and R<sup>8</sup> are selected such that in the resultant compound the carbon atom to which R<sup>7</sup> and R<sup>8</sup> are attached is itself attached to no more than one heteroatom;

R<sup>9</sup> is selected from the group consisting of  
25 hydrogen,  
C<sub>1-8</sub> alkyl,  
aryl,  
aryl C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonyloxy C<sub>1-4</sub> alkyl,  
30 aryl C<sub>1-8</sub> alkylcarbonyloxy C<sub>1-4</sub> alkyl,  
C<sub>1-8</sub> alkylaminocarbonylmethylene, and  
C<sub>1-8</sub> dialkylaminocarbonylmethylene;

R10, R11, R12 and R13 are each independently selected from the group consisting of

- hydrogen,
- C<sub>1-8</sub> alkyl,
- 5 aryl,
- halogen,
- hydroxyl,
- aminocarbonyl,
- C<sub>3-8</sub> cycloalkyl,
- 10 amino C<sub>1-6</sub> alkyl,
- (aryl)<sub>p</sub>aminocarbonyl,
- hydroxycarbonyl,
- (aryl C<sub>1-5</sub> alkyl)<sub>p</sub>aminocarbonyl,
- hydroxycarbonyl C<sub>1-6</sub> alkyl,
- 15 aryl C<sub>1-6</sub> alkyl,
- (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,
- (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,
- C<sub>1-8</sub> alkylsulfonyl,
- C<sub>1-8</sub> alkoxycarbonyl,
- 20 aryloxy carbonyl,
- aryl C<sub>1-8</sub> alkoxycarbonyl,
- C<sub>1-8</sub> alkylcarbonyl,
- arylcarbonyl,
- aryl C<sub>1-6</sub> alkylcarbonyl,
- 25 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,
- aminosulfonyl,
- C<sub>1-8</sub> alkylaminosulfonyl,
- (aryl)<sub>p</sub>aminosulfonyl,
- (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonyl,
- 30 C<sub>1-6</sub> alkylsulfonyl,
- arylsulfonyl,
- aryl C<sub>1-6</sub> alkylsulfonyl,
- aryl C<sub>1-6</sub> alkylcarbonyl,
- C<sub>1-6</sub> alkylthiocarbonyl,

- arylthiocarbonyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonyl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 5 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 10 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 15 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 20 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 25 C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 30 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 35 aryl C<sub>1-6</sub> alkylcarbonyloxy,

- (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
C<sub>1-8</sub> alkylsulfonylamino,  
arylsulfonylamino,  
C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
5 arylsulfonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonylamino,  
aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-8</sub> alkoxycarbonylamino,  
C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
10 aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
aryl C<sub>1-8</sub> alkoxycarbonylamino,  
aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonylamino,  
C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
15 arylcarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonylamino,  
aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
20 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
aminosulfonylamino C<sub>1-6</sub> alkyl,  
25 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
30 C<sub>1-6</sub> alkylsulfonyl,  
C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
arylsulfonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonyl,  
aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,

- C<sub>1-6</sub> alkylcarbonyl,  
C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
arylcarbonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
5 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
10 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl; or  
15 R<sup>10</sup> and R<sup>12</sup> are taken together with the carbon atoms to which  
they are attached to form a 5- to 7-membered monocyclic aromatic  
or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms  
selected from the group consisting of N, O, and S wherein the ring  
nitrogen atoms are unsubstituted or substituted with one R<sup>1</sup>  
20 substituent and the ring carbon atoms are unsubstituted or  
substituted with one or two R<sup>1</sup> substituents,  
and wherein any of the alkyl groups of R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> are either  
unsubstituted or substituted with one to three R<sup>1</sup> substituents;  
25 wherein  
each m is independently an integer from 0 to 6;  
each n is independently an integer from 0 to 6  
each p is independently an integer from 0 to 2;  
each r is independently an integer from 1 to 3;  
30 each s is independently an integer from 0 to 3;  
each t is independently an integer from 0 to 3; and  
each v is independently an integer from 0 to 2;

and the pharmaceutically acceptable salts thereof.

The present invention also relates to pharmaceutical compositions comprising the compounds of the present invention and a pharmaceutically acceptable carrier.

5 The present invention also relates to methods for making the pharmaceutical compositions of the present invention.


The present invention also relates to methods for eliciting an integrin receptor antagonizing effect in a mammal in need thereof by administering the compounds and pharmaceutical compositions of the present invention.

10 The present invention also relates to methods for inhibiting bone resorption, restenosis, atherosclerosis, inflammation, viral disease, diabetic retinopathy, macular degeneration, angiogenesis, wound healing, tumor growth, and metastasis by administering the compounds and pharmaceutical compositions of the present invention.




15 The present invention also relates to methods for treating osteoporosis by administering the compounds and pharmaceutical compositions of the present invention.

#### DETAILED DESCRIPTION OF THE INVENTION



20 The present invention relates to compounds useful as integrin receptor antagonists. Compounds of the present invention are described by the following structural formulas selected from the group consisting of



25




5




, and

10



wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are substituted only with R<sup>10</sup> and R<sup>12</sup>;

**X is selected from the group consisting of**



10 a 5- or 6-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R<sup>1</sup> substituent and the ring carbon atoms are unsubstituted or substituted with one or two R<sup>1</sup> substituents, and

15 a 9- to 14-membered polycyclic ring system, wherein one or more  
of the rings is aromatic, and wherein the polycyclic ring system  
has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of  
N, O, and S wherein the ring nitrogen atoms are unsubstituted or  
substituted with one R<sup>1</sup> substituent and the ring carbon atoms are  
20 unsubstituted or substituted with one or two R<sup>1</sup> substituents;

. Y is selected from the group consisting of

25       $-(\text{CH}_2)_m-$ ,  
           $-(\text{CH}_2)_m-\text{O}-(\text{CH}_2)_n-$ ,  
           $-(\text{CH}_2)_m-\text{NR}^4-(\text{CH}_2)_n-$ .



- 5        $-(CH_2)_m-S-(CH_2)_n-$ ,  
           $-(CH_2)_m-SO-(CH_2)_n-$ ,  
           $-(CH_2)_m-SO_2-(CH_2)_n-$ ,  
           $-(CH_2)_m-O-(CH_2)_n-O-(CH_2)_p-$ ,  
           $-(CH_2)_m-O-(CH_2)_n-NR^4-(CH_2)_p-$ ,  
           $-(CH_2)_m-NR^4-(CH_2)_n-NR^4-(CH_2)_p-$ ,  
           $-(CH_2)_m-O-(CH_2)_n-S-(CH_2)_p-$ ,  
           $-(CH_2)_m-S-(CH_2)_n-S-(CH_2)_p-$ ,  
           $-(CH_2)_m-NR^4-(CH_2)_n-S-(CH_2)_p-$ ,  
 10        $-(CH_2)_m-NR^4-(CH_2)_n-O-(CH_2)_p-$ ,  
           $-(CH_2)_m-S-(CH_2)_n-O-(CH_2)_p-$ ,  
           $-(CH_2)_m-S-(CH_2)_n-NR^4-(CH_2)_p-$ , and  
           $-(CH_2)_m-Z-(CH_2)_n-$ ,

- 15       wherein Z is a 3- to 10-membered monocyclic or polycyclic aromatic or  
          nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected  
          from the group consisting of N, O, and S wherein the 3- to 10-membered  
          monocyclic or polycyclic aromatic or nonaromatic ring system is either  
          unsubstituted or substituted with one or two  $R^1$  substituents, and  
          wherein any methylene ( $CH_2$ ) carbon atom in Y, other than in  $R^4$ , can  
 20       be substituted by one or two  $R^3$  substituents; and

wherein  $R^1$  and  $R^2$  are each independently selected from the group consisting of

- 25       hydrogen, halogen,  $C_{1-10}$  alkyl,  $C_{3-8}$  cycloalkyl,  
           $C_{3-8}$  cycloheteroalkyl,  $C_{3-8}$  cycloalkyl  $C_{1-6}$  alkyl,  
           $C_{3-8}$  cycloheteroalkyl  $C_{1-6}$  alkyl, aryl, aryl  $C_{1-8}$  alkyl, amino,  
          amino  $C_{1-8}$  alkyl,  $C_{1-3}$  acylamino,  $C_{1-3}$  acylamino  $C_{1-8}$  alkyl,  
           $(C_{1-6}$  alkyl) $_p$ amino,  $(C_{1-6}$  alkyl) $_p$ amino  $C_{1-8}$  alkyl,  
           $C_{1-4}$  alkoxy,  $C_{1-4}$  alkoxy  $C_{1-6}$  alkyl, hydroxycarbonyl,  
 30       hydroxycarbonyl  $C_{1-6}$  alkyl,  $C_{1-3}$  alkoxycarbonyl,  
           $C_{1-3}$  alkoxycarbonyl  $C_{1-6}$  alkyl, hydroxycarbonyl-  
           $C_{1-6}$  alkyloxy, hydroxy, hydroxy  $C_{1-6}$  alkyl,  $C_{1-6}$  alkyloxy-  
           $C_{1-6}$  alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,  
          trifluoroethoxy,  $C_{1-8}$  alkyl-S(O) $_p$ ,  $(C_{1-8}$  alkyl) $_p$ aminocarbonyl,

C<sub>1-8</sub> alkyloxycarbonylamino, (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>amino, (aryl)<sub>p</sub>amino, aryl C<sub>1-8</sub>  
 alkylsulfonylamino, and C<sub>1-8</sub> alkylsulfonylamino;  
 or two R<sup>1</sup> substituents, when on the same carbon atom, are taken  
 5 together to form a carbonyl group;

each R<sup>3</sup> is independently selected from the group consisting of  
 hydrogen,  
 aryl,  
 10 C<sub>1-10</sub> alkyl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 15 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 halogen,  
 hydroxyl,  
 oxo,  
 20 trifluoromethyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxy carbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 25 C<sub>1-6</sub> alkylcarbonyloxy,  
 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 amino C<sub>1-6</sub> alkyl,  
 arylaminocarbonyl,  
 30 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 hydroxycarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,

- HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 5 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 10 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 15 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 20 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 25 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 30 C<sub>1-8</sub> alkoxycarbonylamino,  
 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,

- C<sub>1-8</sub> alkylcarbonylamino,  
C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonylamino,  
5 aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
10 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
15 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylsulfonyl,  
C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
20 arylsulfonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonyl,  
aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylcarbonyl,  
C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
25 arylcarbonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
30 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,

(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl;  
or two R<sup>3</sup> substituents, when on the same carbon atom, are taken  
together with the carbon atom to which they are attached to  
5 form a carbonyl group or a cyclopropyl group,  
wherein any of the alkyl groups of R<sup>3</sup> are either unsubstituted or  
substituted with one to three R<sup>1</sup> substituents,  
and provided that each R<sup>3</sup> is selected such that in the resultant  
compound the carbon atom or atoms to which R<sup>3</sup> is attached is itself  
10 attached to no more than one heteroatom;

each R<sup>4</sup> is independently selected from the group consisting of  
hydrogen;  
aryl,  
15 aminocarbonyl,  
C<sub>3-8</sub> cycloalkyl,  
amino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl,  
(aryl C<sub>1-5</sub> alkyl)<sub>p</sub>aminocarbonyl,  
20 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-8</sub> alkyl,  
aryl C<sub>1-6</sub> alkyl,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
(aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
25 C<sub>1-8</sub> alkylsulfonyl,  
C<sub>1-8</sub> alkoxycarbonyl,  
aryloxycarbonyl,  
aryl C<sub>1-8</sub> alkoxycarbonyl,  
C<sub>1-8</sub> alkylcarbonyl,  
30 arylcarbonyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
aminosulfonyl,  
C<sub>1-8</sub> alkylaminosulfonyl,

(aryl)<sub>p</sub>aminosulfonyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonyl,  
 arylsulfonyl,  
 arylC<sub>1-6</sub> alkylsulfonyl,  
 5 C<sub>1-6</sub> alkylthiocarbonyl,  
 arylthiocarbonyl, and  
 aryl C<sub>1-6</sub> alkylthiocarbonyl,  
 wherein any of the alkyl groups of R<sup>4</sup> are either unsubstituted or  
 substituted with one to three R<sup>1</sup> substituents;

10

R<sup>5</sup> and R<sup>6</sup> are each independently selected from the group consisting of  
 hydrogen,  
 C<sub>1-10</sub> alkyl,

15

aryl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 20 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,

20

halogen,  
 hydroxyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,

25

C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,

30

amino C<sub>1-6</sub> alkyl,  
 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,

- hydroxycarbonyl,  
hydroxycarbonyl C<sub>1-6</sub> alkyl,  
HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
5 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
10 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
15 C<sub>1-6</sub> alkoxy,  
aryl C<sub>1-6</sub> alkoxy,  
aryl C<sub>1-6</sub> alkyl,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>amino,  
20 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
(aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
arylcarbonyloxy,  
aryl C<sub>1-6</sub> alkylcarbonyloxy,  
25 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
C<sub>1-8</sub> alkylsulfonylamino,  
arylsulfonylamino,  
C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
arylsulfonylamino C<sub>1-6</sub> alkyl,  
30 aryl C<sub>1-6</sub> alkylsulfonylamino,  
aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-8</sub> alkoxycarbonylamino,  
C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
35 aryl C<sub>1-8</sub> alkoxycarbonylamino,

- aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonylamino,  
C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcabonylamino C<sub>1-6</sub> alkyl,  
5 aryl C<sub>1-6</sub> alkylcarbonylamino,  
aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
10 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
15 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylsulfonyl,  
20 C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
arylsulfonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonyl,  
aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylcarbonyl,  
25 C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
arylcabonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
30 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,



(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl;

or R<sup>5</sup> and R<sup>6</sup> are taken together with the carbon atom to which  
 5 they are attached to form a carbonyl group,  
 wherein any of the alkyl groups of R<sup>5</sup> or R<sup>6</sup> are either unsubstituted or  
 substituted with one to three R<sup>1</sup> substituents,  
 and provided that each R<sup>5</sup> and R<sup>6</sup> are selected such that in the resultant  
 compound the carbon atom to which R<sup>5</sup> and R<sup>6</sup> are attached is itself  
 10 attached to no more than one heteroatom;

R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of  
 hydrogen,  
 C<sub>1-10</sub> alkyl,  
 15 aryl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 20 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 halogen,  
 hydroxyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 25 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 C<sub>3-8</sub> cycloalkyl,  
 30 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 amino C<sub>1-6</sub> alkyl,  
 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 aminocarbonyl,

- aminocarbonyl C<sub>1-6</sub> alkyl,  
 hydroxycarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 5 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 10 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 15 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 20 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 25 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 30 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkoxycarbonylamino,  
 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,

- aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
aryl C<sub>1-8</sub> alkoxycarbonylamino,  
aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonylamino,  
5 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonylamino,  
aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
10 (C<sub>1-8</sub> alkyl)paminocarbonylamino,  
(C<sub>1-8</sub> alkyl)paminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl)paminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)paminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)paminocarbonylamino C<sub>1-6</sub> alkyl,  
15 aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)paminosulfonylamino,  
(C<sub>1-8</sub> alkyl)paminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl)paminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)paminosulfonylamino,  
20 (aryl C<sub>1-8</sub> alkyl)paminosulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylsulfonyl,  
C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
arylsulfonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonyl,  
25 aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylcarbonyl,  
C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
arylcarbonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
30 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,

- aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
5 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl, and  
C<sub>7-20</sub> polycyclyl C<sub>0-8</sub> alkylsulfonylamino;

wherein any of the alkyl groups of R<sup>7</sup> and R<sup>8</sup> are either unsubstituted or substituted with one to three R<sup>1</sup> substituents,  
and provided that each R<sup>7</sup> and R<sup>8</sup> are selected such that in the resultant  
10 compound the carbon atom to which R<sup>7</sup> and R<sup>8</sup> are attached is itself  
attached to no more than one heteroatom;

- R<sup>9</sup> is selected from the group consisting of  
hydrogen,  
15 C<sub>1-8</sub> alkyl,  
aryl,  
aryl C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonyloxy C<sub>1-4</sub> alkyl,  
aryl C<sub>1-8</sub> alkylcarbonyloxy C<sub>1-4</sub> alkyl,  
20 C<sub>1-8</sub> alkylaminocarbonylmethylene, and  
C<sub>1-8</sub> dialkylaminocarbonylmethylene;

- R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup> and R<sup>13</sup> are each independently selected from the group  
consisting of  
25 hydrogen,  
C<sub>1-8</sub> alkyl,  
aryl,  
halogen,  
hydroxyl,  
30 aminocarbonyl,  
C<sub>3-8</sub> cycloalkyl,  
amino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminocarbonyl,  
hydroxycarbonyl,

- (aryl C<sub>1-5</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 5 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
 C<sub>1-8</sub> alkylsulfonyl,  
 C<sub>1-8</sub> alkoxycarbonyl,  
 aryloxycarbonyl,  
 aryl C<sub>1-8</sub> alkoxycarbonyl,  
 10 C<sub>1-8</sub> alkylcarbonyl,  
 arylcarbonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 aminosulfonyl,  
 15 C<sub>1-8</sub> alkylaminosulfonyl,  
 (aryl)<sub>p</sub>aminosulfonyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonyl,  
 C<sub>1-6</sub> alkylsulfonyl,  
 arylsulfonyl,  
 20 aryl C<sub>1-6</sub> alkylsulfonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl,  
 C<sub>1-6</sub> alkylthiocarbonyl,  
 arylthiocarbonyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonyl,  
 25 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 30 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,

- C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 5 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 10 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 15 aminocarbonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 20 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 25 C<sub>1-8</sub> alkylsulfonylamino,  
 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 30 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkoxycarbonylamino,  
 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino,

- aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonylamino,  
C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcarbonylamino C<sub>1-6</sub> alkyl,  
5 aryl C<sub>1-6</sub> alkylcarbonylamino,  
aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
10 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
15 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylsulfonyl,  
20 C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
arylsulfonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonyl,  
aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylcarbonyl,  
25 C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
arylcarbonyl C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonyl,  
aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
C<sub>1-6</sub> alkylthiocarbonylamino,  
30 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,

(aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl; or


5 R<sup>10</sup> and R<sup>12</sup> are taken together with the carbon atoms to which  
 they are attached to form a 5- to 7-membered monocyclic aromatic  
 or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms  
 selected from the group consisting of N, O, and S wherein the ring  
 nitrogen atoms are unsubstituted or substituted with one R<sup>1</sup>  
 substituent and the ring carbon atoms are unsubstituted or  
 10 substituted with one or two R<sup>1</sup> substituents,  
 and wherein any of the alkyl groups of R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> are either  
 unsubstituted or substituted with one to three R<sup>1</sup> substituents;

wherein

15 each m is independently an integer from 0 to 6;  
 each n is independently an integer from 0 to 6  
 each p is independently an integer from 0 to 2;  
 each r is independently an integer from 1 to 3;  
 each s is independently an integer from 0 to 3;  
 20 each t is independently an integer from 0 to 3; and  
 each v is independently an integer from 0 to 2;


and the pharmaceutically acceptable salts thereof.

25 In one embodiment of the present invention, compounds are  
 described by the following structural formulas selected from the group  
 consisting of




30






and



wherein the dotted line a represents a single or a double bond, provided  
 5 that when a represents a double bond, the double bond carbon atoms are  
 substituted only with R<sup>10</sup> and R<sup>12</sup>.


In a class of this embodiment of the present invention,  
 compounds are described by the following structural formula

10



wherein the dotted line a represents a single or a double bond, provided  
 that when a represents a double bond, the double bond carbon atoms are  
 15 substituted only with R<sup>10</sup> and R<sup>12</sup>.

In a subclass of this class of the present invention,  
 compounds are described by the following structural formula




In the compounds of the present invention, X is preferably  
 a 6-membered monocyclic aromatic ring system having 1 or 2 nitrogen  
 5 atoms wherein each ring carbon atom is unsubstituted or substituted  
 with one R<sup>1</sup> substituent, or

a 9- to 14-membered polycyclic ring system, wherein one or more of the  
 rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3  
 10 or 4 heteroatoms selected from the group consisting of N, O, and S  
 wherein the ring nitrogen atoms are unsubstituted or substituted with  
 one R<sup>1</sup> substituent and the ring carbon atoms are unsubstituted or  
 substituted with one or two R<sup>1</sup> substituents.


15

More preferably, X is selected from the group consisting of



20

Most preferably X is




In the compounds of the present invention, Y is preferably  
5 selected from the group consisting of

- (CH<sub>2</sub>)<sub>m</sub>-,
- (CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-,
- (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-,
- 10 -(CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>-,
- (CH<sub>2</sub>)<sub>m</sub>-SO-(CH<sub>2</sub>)<sub>n</sub>-,
- (CH<sub>2</sub>)<sub>m</sub>-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>n</sub>-,
- (CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,
- (CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-,
- 15 -(CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-, and
- (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,

wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than in R<sup>4</sup>, can be substituted by one or two R<sup>3</sup> substituents.

20 More preferably Y is selected from the group consisting of



wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than in R<sup>4</sup>, can  
25 be substituted by one or two R<sup>3</sup> substituents.

Most preferably Y is (CH<sub>2</sub>)<sub>m</sub> or (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>  
wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than R<sup>4</sup>, can be  
substituted by one or two R<sup>3</sup> substituents.

30

In the compounds of the present invention, R<sup>1</sup> and R<sup>2</sup> are preferably selected from the group consisting of hydrogen, halogen, C<sub>1</sub>-10 alkyl, C<sub>3</sub>-8 cycloalkyl, C<sub>3</sub>-8 cycloheteroalkyl, hydroxy, nitro, cyano, trifluoromethyl, and trifluoromethoxy.

More preferably, R<sup>1</sup> and R<sup>2</sup> are selected from the group consisting of hydrogen, halogen, C<sub>1</sub>-10 alkyl, C<sub>3</sub>-8 cycloalkyl, trifluoromethyl, and trifluoromethoxy.

In the compounds of the present invention, R<sup>3</sup> is preferably selected from the group consisting of

hydrogen,  
fluoro,  
trifluoromethyl,  
aryl,  
C<sub>1</sub>-8 alkyl,  
arylC<sub>1</sub>-6 alkyl  
hydroxyl,  
oxo,  
arylamino carbonyl,  
aryl C<sub>1</sub>-5 alkylamino carbonyl,  
amino carbonyl, and  
amino carbonyl C<sub>1</sub>-6 alkyl.

More preferably, R<sup>3</sup> is selected from the group consisting of  
fluoro,  
aryl,  
C<sub>1</sub>-8 alkyl,  
arylC<sub>1</sub>-6 alkyl  
hydroxyl,  
oxo, and  
arylamino carbonyl.

In the compounds of the present invention, R<sup>4</sup> is preferably selected from the group consisting of

hydrogen,  
aryl,  
C<sub>3-8</sub> cycloalkyl,  
C<sub>1-8</sub> alkyl,  
5 C<sub>1-8</sub> alkylcarbonyl,  
arylcarbonyl,  
C<sub>1-6</sub> alkylsulfonyl,  
arylsulfonyl,  
arylC<sub>1-6</sub>alkylsulfonyl,  
10 arylC<sub>1-6</sub>alkylcarbonyl,  
C<sub>1-8</sub>alkylaminocarbonyl,  
arylC<sub>1-5</sub>alkylaminocarbonyl,  
arylC<sub>1-8</sub>alkoxycarbonyl, and  
C<sub>1-8</sub>alkoxycarbonyl.

15

More preferably, R<sup>4</sup> is selected from the group consisting of  
hydrogen,  
C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonyl,  
20 arylcarbonyl,  
arylC<sub>1-6</sub>alkylcarbonyl,  
C<sub>1-6</sub> alkylsulfonyl,  
arylsulfonyl, and  
arylC<sub>1-6</sub>alkylsulfonyl.

25

In one embodiment of the present invention, R<sup>5</sup> and R<sup>6</sup> are  
each independently selected from the group consisting of  
hydrogen,  
aryl,  
30 C<sub>1-8</sub> alkyl,  
aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
aryl C<sub>1-6</sub> alkyl,  
CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-, and  
HC≡C-(CH<sub>2</sub>)<sub>t</sub>-.

In a class of this embodiment of the present invention,  $R^6$  is hydrogen and  $R^5$  is selected from the group consisting of

hydrogen,  
 aryl,  
 5 C<sub>1-8</sub> alkyl,  
 aryl-C $\equiv$ C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl C<sub>1-6</sub> alkyl,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-, and  
 HC $\equiv$ C-(CH<sub>2</sub>)<sub>t</sub>-.

10 In a subclass of this class of the present invention,  $R^6$ ,  $R^7$ , and  $R^8$  are each hydrogen and  $R^5$  is selected from the group consisting of

hydrogen,  
 aryl,  
 15 C<sub>1-8</sub> alkyl,  
 aryl-C $\equiv$ C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl C<sub>1-6</sub> alkyl,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-, and  
 HC $\equiv$ C-(CH<sub>2</sub>)<sub>t</sub>-.

20 In another embodiment of the present invention,  $R^7$  and  $R^8$  are each independently selected from the group consisting of

hydrogen,  
 aryl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 25 arylcarbonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 30 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkoxy carbonylamino,  
 C<sub>1-8</sub> alkoxy carbonylamino C<sub>1-8</sub> alkyl,  
 aryloxy carbonylamino C<sub>1-8</sub> alkyl,

- aryl C<sub>1-8</sub> alkoxy-carbonylamino,  
 aryl C<sub>1-8</sub> alkoxy-carbonylamino C<sub>1-8</sub> alkyl,  
 C<sub>1-8</sub> alkyl-carbonylamino C<sub>1-6</sub> alkyl,  
 aryl-carbonylamino C<sub>1-6</sub> alkyl,  
 5 aryl C<sub>1-6</sub> alkyl-carbonylamino,  
 aryl C<sub>1-6</sub> alkyl-carbonylamino C<sub>1-6</sub> alkyl,  
 aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 10 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 15 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylthiocarbonylamino,  
 20 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino, and  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl.

- 25 In a class of this embodiment of the present invention, R<sup>8</sup> is  
 hydrogen and R<sup>7</sup> is selected from the group consisting of consisting of  
 hydrogen,  
 aryl,  
 C<sub>1-8</sub> alkyl-carbonylamino,  
 30 aryl C<sub>1-6</sub> alkyl-carbonylamino,  
 aryl-carbonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 arylsulfonylamino,

C<sub>1-8</sub> alkoxy carbonylamino,  
aryl C<sub>1-8</sub> alkoxy carbonylamino,  
arylamino carbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub> aminocarbonylamino,  
5 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub> aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub> aminosulfonylamino, and  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub> aminosulfonylamino.

In a subclass of this class of the present invention, R<sup>5</sup>, R<sup>6</sup>,  
and R<sup>8</sup> are each hydrogen and R<sup>7</sup> is selected from the group consisting  
10 of

hydrogen,  
aryl,  
C<sub>1-8</sub> alkyl carbonylamino,  
aryl C<sub>1-6</sub> alkyl carbonylamino,  
15 aryl carbonylamino,  
C<sub>1-8</sub> alkyl sulfonylamino,  
aryl C<sub>1-6</sub> alkyl sulfonylamino,  
aryl sulfonylamino,  
C<sub>1-8</sub> alkoxy carbonylamino,  
20 aryl C<sub>1-8</sub> alkoxy carbonylamino,  
arylamino carbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub> aminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub> aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub> aminosulfonylamino, and  
25 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub> aminosulfonylamino.

In the compounds of the present invention, R<sup>9</sup> is preferably  
selected from the group consisting of hydrogen, methyl, and ethyl.

More preferably, R<sup>9</sup> is hydrogen.

30 In the compounds of the present invention, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>,  
and R<sup>13</sup> are preferably each independently selected from the group  
consisting of hydrogen, aryl, C<sub>1-6</sub> alkyl, and aryl C<sub>1-6</sub> alkyl.

In the compounds of the present invention, m is preferably  
an integer from 0 to 4, and more preferably from 0 to 3.



In the compounds of the present invention, n is preferably an integer from 0 to 4, more preferably from 0 to 3.



In the compounds of the present invention, r is preferably an integer from 1 to 2.

5 In the compounds of the present invention, s is preferably an integer from 0 to 2.

In the compounds of the present invention, t is preferably an integer from 0 to 2, more preferably from 0 to 1.


In the compounds of the present invention, v is preferably 0.

10 In certain embodiments of the present invention, the compounds correspond to the formulas with the following designated stereochemistry at the carbon atom where R<sup>5</sup> and R<sup>6</sup> are attached:



15

, and



;

wherein the substituents X, Y, Z, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> and the subscripts a, m, n, p, r, s, t, and v are as described above.

5 Illustrative but nonlimiting examples of compounds of the present invention that are useful as integrin receptor antagonists are the following:

- 10 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 20 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 25 3-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,  
trifluoroacetate salt,
- 3(S)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,  
bis(trifluoroacetate) salt,
- 3(R)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
25 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-  
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-  
35 propionic acid,

- 3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 5 3-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 10 3(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 15 3-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 20 3(R)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 30 3(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 35

- 3(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 5 3(R)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 10 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 20 3(S)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 2(S)-Benzenesulfonylamino-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
30 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-  
imidazolidin-1-yl)-pent-4-enoic acid,

- 3(R)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 5 3-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 10 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 15 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(R)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 20 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 25 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 30 3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 35

- 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10 3-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,



3-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

5 3-(Furo[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

10 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

3(R)-(Furo[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

15 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

20 3-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

25 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

3(R)-(Furo[3,2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

30 3-(Furo[3,2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

- 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 15 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
20 [1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-  
yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
35 [1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

5 3-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

10 Further illustrative of the present invention are the compounds selected from the group consisting of

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15

3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20 3(S)-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25 3(S)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

30

3(S)-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

35 3(R)-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 10 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3(R)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(S)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid, bis(trifluoroacetate) salt,
- 30 3(R)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(R)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3(S)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(R)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(R)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35 3(S)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(R)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-  
15 imidazolidin-1-yl}-pent-4-enoic acid,
- 3(R)-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-  
imidazolidin-1-yl}-pent-4-enoic acid,
- 20 3(S)-(5-Ethoxy-pyridin-3-yl)-3-{3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-  
yl]-propyl]-2-oxo-imidazolidin-1-yl}-propionic acid,
- 3(R)-(5-Ethoxy-pyridin-3-yl)-3-{3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-  
yl]-propyl]-2-oxo-imidazolidin-1-yl}-propionic acid,
- 25 3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3(S)-(5-  
ethoxy-pyridin-3-yl)-propionic acid,
- 3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3(R)-(5-  
30 ethoxy-pyridin-3-yl)-propionic acid,
- 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-  
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-  
yl}propionic acid,
- 35

- 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 15 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 20 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
35 2-yl)propyl]imidazolidin-1-yl)propionic acid,



3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

5 3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

10 Yet further illustrative are the compounds

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15 3(S)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20

3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25 3(S)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,  
bis(trifluoroacetate) salt,

30

3(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

35 3(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

5 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

10 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

15

For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts.

20 Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid. Representative salts include the following: acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium, camsylate, carbonate, chloride, clavulanate, citrate, 25 dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, 30 napsylate, nitrate, N-methylglucamine ammonium salt, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide and valerate.

35 Furthermore, where the compounds of the invention carry an acidic

moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts.

5           The compounds of the present invention can have chiral centers and occur as racemates, racemic mixtures, diastereomeric mixtures, and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention. Therefore, where a compound is chiral, the separate enantiomers or  
10       diastereomers, substantially free of the other, are included within the scope of the invention; further included are all mixtures of the two enantiomers. Also included within the scope of the invention are polymorphs and hydrates of the compounds of the instant invention.

          The present invention includes within its scope prodrugs of  
15       the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with  
20       the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," ed. H. Bundgaard, Elsevier, 1985,  
25       which is incorporated by reference herein in its entirety. Metabolites of these compounds include active species produced upon introduction of compounds of this invention into the biological milieu.

          The term "therapeutically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the  
30       biological or medical response of a tissue, system, animal or human that is being sought by a researcher or clinician.

          The term "integrin receptor antagonist," as used herein, refers to a compound which binds to and antagonizes either the  $\alpha\beta 3$  receptor, the  $\alpha\beta 5$  receptor, or the  $\alpha\beta 6$  receptor, or a compound which

binds to and antagonizes combinations of these receptors (for example, a dual  $\alpha\beta 3/\alpha\beta 5$  receptor antagonist).

The term "bone resorption," as used herein, refers to the process by which osteoclasts degrade bone.

5       The term "alkyl" shall mean straight or branched chain alkanes of one to ten total carbon atoms, or any number within this range (i.e., methyl, ethyl, 1-propyl, 2-propyl, n-butyl, s-butyl, t-butyl, etc.).

10       The term "alkenyl" shall mean straight or branched chain alkenes of two to ten total carbon atoms, or any number within this range.

The term "alkynyl" shall mean straight or branched chain alkynes of two to ten total carbon atoms, or any number within this range.

15       The term "cycloalkyl" shall mean cyclic rings of alkanes of three to eight total carbon atoms, or any number within this range (i.e., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl).

20       The term "cycloheteroalkyl," as used herein, shall mean a 3- to 8-membered fully saturated heterocyclic ring containing one or two heteroatoms chosen from N, O or S. Examples of cycloheteroalkyl groups include, but are not limited to piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, piperazinyl.

25       The term "alkoxy," as used herein, refers to straight or branched chain alkoxides of the number of carbon atoms specified (e.g., C<sub>1-5</sub> alkoxy), or any number within this range (i.e., methoxy, ethoxy, etc.).

30       The term "aryl," as used herein, refers to a monocyclic or polycyclic system comprising at least one aromatic ring, wherein the monocyclic or polycyclic system contains 0, 1, 2, 3, or 4 heteroatoms chosen from N, O, or S, and wherein the monocyclic or polycyclic system is either unsubstituted or substituted with one or more groups independently selected from hydrogen, halogen, C<sub>1-10</sub> alkyl, C<sub>3-8</sub> cycloalkyl, aryl, aryl C<sub>1-8</sub> alkyl, amino, amino C<sub>1-8</sub> alkyl, C<sub>1-3</sub> acylamino, C<sub>1-3</sub> acylamino C<sub>1-8</sub> alkyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkylamino C<sub>1-8</sub> alkyl, C<sub>1-6</sub> dialkylamino, C<sub>1-6</sub> dialkylamino-C<sub>1-8</sub> alkyl, C<sub>1-4</sub> alkoxy,

35

C<sub>1-4</sub> alkoxy C<sub>1-6</sub> alkyl, hydroxycarbonyl, hydroxycarbonyl C<sub>1-6</sub> alkyl, C<sub>1-5</sub> alkoxycarbonyl, C<sub>1-3</sub> alkoxycarbonyl C<sub>1-6</sub> alkyl, hydroxycarbonyl C<sub>1-6</sub> alkyloxy, hydroxy, hydroxy C<sub>1-6</sub> alkyl, cyano, trifluoromethyl, oxo or C<sub>1-5</sub> alkylcarbonyloxy. Examples of aryl include, but are not limited to,

5 phenyl, naphthyl, pyridyl, pyrazinyl, pyrimidinyl, imidazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, indolyl, thienyl, furyl, pyrrolyl, pyrazolyl, dihydrobenzofuryl, benzo(1,3) dioxolane, oxazolyl, isoxazolyl and thiazolyl, which are either unsubstituted or substituted with one or more groups independently selected from hydrogen,

10 halogen, C<sub>1-10</sub> alkyl, C<sub>3-8</sub> cycloalkyl, aryl, aryl C<sub>1-8</sub> alkyl, amino, amino C<sub>1-8</sub> alkyl, C<sub>1-3</sub> acylamino, C<sub>1-3</sub> acylamino C<sub>1-8</sub> alkyl, C<sub>1-6</sub> alkylamino, C<sub>1-6</sub> alkylamino-C<sub>1-8</sub> alkyl, C<sub>1-6</sub> dialkylamino, C<sub>1-6</sub> dialkylamino C<sub>1-8</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkoxy C<sub>1-6</sub> alkyl, hydroxycarbonyl, hydroxycarbonyl C<sub>1-6</sub> alkyl, C<sub>1-5</sub> alkoxycarbonyl, C<sub>1-3</sub> alkoxycarbonyl

15 C<sub>1-6</sub> alkyl, hydroxycarbonyl C<sub>1-6</sub> alkyloxy, hydroxy, hydroxy C<sub>1-6</sub> alkyl, cyano, trifluoromethyl, oxo or C<sub>1-5</sub> alkylcarbonyloxy. Preferably, the aryl group is unsubstituted, mono-, di-, tri- or tetra-substituted with one to four of the above-named substituents; more preferably, the aryl group is unsubstituted, mono-, di- or tri-substituted with one to three of the

20 above-named substituents; most preferably, the aryl group is unsubstituted, mono- or di-substituted with one to two of the above-named substituents.

Whenever the term "alkyl" or "aryl" or either of their prefix roots appears in a name of a substituent (e.g., aryl C<sub>0-8</sub> alkyl), it shall be

25 interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C<sub>1-10</sub>) shall refer independently to the number of carbon atoms in an alkyl or cyclic alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

30 The terms "arylalkyl" and "alkylaryl" include an alkyl portion where alkyl is as defined above and to include an aryl portion where aryl is as defined above. Examples of arylalkyl include, but are not limited to, benzyl, fluorobenzyl, chlorobenzyl, phenylethyl, phenylpropyl, fluorophenylethyl, chlorophenylethyl, thienylmethyl,

thienylethyl, and thienylpropyl. Examples of alkylaryl include, but are not limited to, toluene, ethylbenzene, propylbenzene, methylpyridine, ethylpyridine, propylpyridine and butylpyridine.

In the compounds of the present invention, two R<sup>1</sup> substituents, when on the same carbon atom, can be taken together with the carbon to which they are attached to form a carbonyl group.

In the compounds of the present invention, two R<sup>3</sup> substituents, when on the same carbon atom, can be taken together with the carbon atom to which they are attached to form a carbonyl group. In such instances, the limitation, that in the resultant compound the carbon atom or atoms to which R<sup>3</sup> is attached is itself attached to no more than one heteroatom, does not apply. Also, in the compounds of the present invention, two R<sup>3</sup> substituents, when on the same carbon atom, can be taken together with the carbon atom to which they are attached to form a cyclopropyl group.

In the compounds of the present invention, R<sup>5</sup> and R<sup>6</sup> can be taken together with the carbon atom to which they are attached to form a carbonyl group. In such instances, the limitation, that in the resultant compound the carbon atom at which R<sup>5</sup> and R<sup>6</sup> is attached is itself attached to no more than one heteroatom, does not apply.

When substituents R<sup>7</sup> and R<sup>8</sup> include the definition C<sub>0</sub> (e.g., C<sub>0-8</sub> alkyl), the group modified by C<sub>0</sub> is not present in the substituent when C is zero. Similarly, when any of the variables m, n, t, or v, is zero, then the group modified by the variable is not present; for example, when t is zero, the group  $-(CH_2)_tC\equiv CH$  is  $-C\equiv CH$ . In addition, the substituent "(C<sub>1-6</sub> alkyl)<sub>p</sub>amino" where p is zero, one or two, refers to an amino, C<sub>1-6</sub> alkylamino and C<sub>1-6</sub> dialkylamino group, respectively. When a C<sub>1-6</sub> dialkylamino substituent is intended, the C<sub>1-6</sub> alkyl groups can be the same (e.g., dimethylamino) or different (e.g., N(CH<sub>3</sub>)(CH<sub>2</sub>CH<sub>3</sub>)). Similarly, the substituent "(aryl)<sub>p</sub>amino" or ["(aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino"], where p is zero, one or two, refers to an amino, arylamino and diarylamino group, [or an amino, aryl C<sub>1-6</sub> alkylamino or di-(aryl C<sub>1-6</sub> alkyl)amino] respectively, where the aryl [or aryl C<sub>1-6</sub> alkyl] groups in a diarylamino [or di-(aryl C<sub>1-6</sub> alkyl)amino] substituent can be the same or different.


In the compounds of the present invention, R<sup>10</sup> and R<sup>12</sup> can be taken together with the carbon atoms to which they are attached to form a 5- to 7-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of N, O, and S wherein said 5- to 7-membered monocyclic aromatic or nonaromatic ring system is either unsubstituted or substituted with one or more R<sup>1</sup> substituents.

The term "halogen" shall include iodine, bromine, chlorine, and fluorine.

The term "oxy" means an oxygen (O) atom. The term "thio" means a sulfur (S) atom. The term "oxo" means "=O." The term "carbonyl" means "C=O."

The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently substituted, it is meant that the (two or more) substituents can be the same or different.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. For example, a C<sub>1-5</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl substituent is equivalent to



In choosing compounds of the present invention, one of ordinary skill in the art will recognize that the various substituents, i.e. X, Y, Z, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> and the subscripts m, n, p, r, s, t, and v are to be chosen in conformity with well-known principles of chemical structure connectivity.

Representative compounds of the present invention typically display submicromolar affinity for the integrin receptors, particularly

the  $\alpha v\beta 3$ ,  $\alpha v\beta 5$ , and/or  $\alpha v\beta 6$  receptors. Compounds of this invention are therefore useful for treating mammals suffering from a bone condition caused or mediated by increased bone resorption, who are in need of such therapy. Pharmacologically effective amounts of the compounds, including pharmaceutically acceptable salts thereof, are administered to the mammal, to inhibit the activity of mammalian osteoclasts.

The compounds of the present invention are administered in dosages effective to antagonize the  $\alpha v\beta 3$  receptor where such treatment is needed, as, for example, in the prevention or treatment of osteoporosis.

Further exemplifying the invention is the method wherein the integrin receptor antagonizing effect is an  $\alpha v\beta 3$  antagonizing effect. An illustration of the invention is the method wherein the  $\alpha v\beta 3$  antagonizing effect is selected from inhibition of bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, tumor growth, or metastasis. Preferably, the  $\alpha v\beta 3$  antagonizing effect is the inhibition of bone resorption.

An example of the invention is the method wherein the integrin receptor antagonizing effect is an  $\alpha v\beta 5$  antagonizing effect. More specifically, the  $\alpha v\beta 5$  antagonizing effect is selected from inhibition of restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, tumor growth, or metastasis.

Illustrating the invention is the method wherein the integrin receptor antagonizing effect is a dual  $\alpha v\beta 3/\alpha v\beta 5$  antagonizing effect. More particularly, the dual  $\alpha v\beta 3/\alpha v\beta 5$  antagonizing effect is selected from inhibition of bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, tumor growth, or metastasis.

Illustrating the invention is the method wherein the integrin receptor antagonizing effect is an  $\alpha v\beta 6$  antagonizing effect. More particularly, the  $\alpha v\beta 6$  antagonizing effect is selected from inhibition of angiogenesis, inflammatory response, or wound healing.

Illustrating the invention is the method wherein the  $\alpha v\beta 3$  antagonizing effect is selected from inhibition of bone resorption,



inhibition of restenosis, inhibition of angiogenesis, inhibition of diabetic retinopathy, inhibition of macular degeneration, inhibition of atherosclerosis, inflammation, viral disease, or inhibition of tumor growth or metastasis. Preferably, the  $\alpha v \beta 3$  antagonizing effect is the inhibition of bone resorption.

More particularly illustrating the invention is a pharmaceutical composition comprising any of the compounds described above and a pharmaceutically acceptable carrier. Another example of the invention is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. Another illustration of the invention is a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

Further illustrating the invention is a method of treating and/or preventing a condition mediated by antagonism of an integrin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds described above. Preferably, the condition is selected from bone resorption, osteoporosis, restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, viral disease, cancer, tumor growth, and metastasis. More preferably, the condition is selected from osteoporosis and cancer. Most preferably, the condition is osteoporosis.

More specifically exemplifying the invention is a method of eliciting an integrin antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above. Preferably, the integrin antagonizing effect is an  $\alpha v \beta 3$  antagonizing effect; more specifically, the  $\alpha v \beta 3$  antagonizing effect is selected from inhibition of bone resorption, inhibition of restenosis, inhibition of atherosclerosis, inhibition of angiogenesis, inhibition of diabetic retinopathy, inhibition of macular degeneration, inhibition of inflammation, inhibition of viral disease, or inhibition of tumor growth or metastasis. Most preferably, the  $\alpha v \beta 3$

antagonizing effect is inhibition of bone resorption. Alternatively, the integrin antagonizing effect is an  $\alpha v \beta 5$  antagonizing effect, an  $\alpha v \beta 6$  antagonizing effect, or a mixed  $\alpha v \beta 3$ ,  $\alpha v \beta 5$ , and  $\alpha v \beta 6$  antagonizing effect. Examples of  $\alpha v \beta 5$  antagonizing effects are inhibition of

5 restenosis, atherosclerosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, or tumor growth. Examples of dual  $\alpha v \beta 6$  antagonizing effects are inhibition of angiogenesis, inflammatory response and wound healing.

Additional examples of the invention are methods of  
10 inhibiting bone resorption and of treating and/or preventing osteoporosis in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above.

Additional illustrations of the invention are methods of  
15 treating hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, periarticular erosions in rheumatoid arthritis, Paget's disease, immobilization-induced osteopenia, and glucocorticoid treatment in a mammal in need thereof, comprising administering to the mammal a therapeutically  
20 effective amount of any of the compounds or any of the pharmaceutical compositions described above.

More particularly exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment and/or prevention of osteoporosis in a  
25 mammal in need thereof. Still further exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment and/or prevention of bone resorption, tumor growth, cancer, restenosis, atherosclerosis, diabetic retinopathy, macular degeneration, inflammation, viral disease, and/or  
30 angiogenesis.

Also exemplifying the invention are compositions further comprising an active ingredient selected from the group consisting of

- a.) an organic bisphosphonate or a pharmaceutically acceptable salt or ester thereof,
- 35 b.) an estrogen receptor modulator,

- c.) a cytotoxic/antiproliferative agent,
- d.) a matrix metalloproteinase inhibitor,
- e.) an inhibitor of epidermal-derived, fibroblast-derived, or platelet-derived growth factors,
- 5 f.) an inhibitor of VEGF,
- g.) an inhibitor of Flk-1/KDR, Flt-1, Tck/Tie-2, or Tie-1,
- h.) a cathepsin K inhibitor; and
- i.) a prenylation inhibitor, such as a farnesyl transferase inhibitor or a geranylgeranyl transferase inhibitor or a dual
- 10 farnesyl/geranylgeranyl transferase inhibitor; and mixtures thereof.

(See, B. Millauer et al., "Dominant-Negative Inhibition of Flk-1 Suppresses the Growth of Many Tumor Types in *Vivo*", Cancer Research, 56, 1615-1620 (1996), which is incorporated by reference herein  
15 in its entirety).

Preferably, the active ingredient is selected from the group consisting of:

- a.) an organic bisphosphonate or a pharmaceutically acceptable salt or ester thereof,
- 20 b.) an estrogen receptor modulator, and
- c.) a cathepsin K inhibitor; and mixtures thereof.

Nonlimiting examples of such bisphosphonates include alendronate, etidronate, pamidronate, risedronate, ibandronate, and pharmaceutically acceptable salts and esters thereof. A particularly  
25 preferred bisphosphonate is alendronate, especially alendronate monosodium trihydrate.

Nonlimiting examples of estrogen receptor modulators include estrogen, progesterin, estradiol, droloxifene, raloxifene, and tamoxifene.

30 Nonlimiting examples of cytotoxic/antiproliferative agents are taxol, vincristine, vinblastine, and doxorubicin.

Cathepsin K, formerly known as cathepsin O2, is a cysteine protease and is described in PCT International Application Publication No. WO 96/13523, published May 9, 1996; U.S. Patent No. 5,501,969,  
35 issued March 3, 1996; and U.S. Patent No. 5,736,357, issued April 7, 1998,

all of which are incorporated by reference herein in their entirety.

Cysteine proteases, specifically cathepsins, are linked to a number of disease conditions, such as tumor metastasis, inflammation, arthritis, and bone remodeling. At acidic pH's, cathepsins can degrade type-I

5 collagen. Cathepsin protease inhibitors can inhibit osteoclastic bone resorption by inhibiting the degradation of collagen fibers and are thus useful in the treatment of bone resorption diseases, such as osteoporosis.

The present invention is also directed to combinations of the compounds of the present invention with one or more agents useful in  
10 the prevention or treatment of osteoporosis. For example, the compounds of the instant invention may be effectively administered in combination with effective amounts of other agents such as an organic bisphosphonate, an estrogen receptor modulator, or a cathepsin K inhibitor.

15 Additional illustrations of the invention are methods of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound described above and one or more agents known to be cytotoxic/antiproliferative. Also, the compounds of the present invention  
20 can be administered in combination with radiation therapy for treating tumor growth and metastasis.

In addition, the integrin  $\alpha v \beta 3$  antagonist compounds of the present invention may be effectively administered in combination with a growth hormone secretagogue in the therapeutic or prophylactic  
25 treatment of disorders in calcium or phosphate metabolism and associated diseases. These diseases include conditions which can benefit from a reduction in bone resorption. A reduction in bone resorption should improve the balance between resorption and formation, reduce bone loss or result in bone augmentation. A reduction  
30 in bone resorption can alleviate the pain associated with osteolytic lesions and reduce the incidence and/or growth of those lesions. These diseases include: osteoporosis (including estrogen deficiency, immobilization, glucocorticoid induced and senile), osteodystrophy, Paget's disease, myositis ossificans, Bechterew's disease, malignant  
35 hypercalcemia, metastatic bone disease, periodontal disease,

cholelithiasis, nephrolithiasis, urolithiasis, urinary calculus, hardening of the arteries (sclerosis), arthritis, bursitis, neuritis and tetany. Increased bone resorption can be accompanied by pathologically high calcium and phosphate concentrations in the plasma, which would  
5 be alleviated by this treatment. Similarly, the present invention would be useful in increasing bone mass in patients with growth hormone deficiency. Thus, preferred combinations are simultaneous or alternating treatments of an  $\alpha v \beta 3$  receptor antagonist of the present invention and a growth hormone secretagogue, optionally including a  
10 third component comprising an organic bisphosphonate, preferably alendronate monosodium trihydrate.

In accordance with the method of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or  
15 concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment, and the term "administering" is to be interpreted accordingly. It will be understood that the scope of combinations of the compounds of this invention with other agents  
20 useful for treating integrin-mediated conditions includes in principle any combination with any pharmaceutical composition useful for treating osteoporosis.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the  
25 specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of  
30 which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups and emulsions. Likewise, they may also be administered in intravenous (bolus or infusion), intraperitoneal, topical (e.g., ocular eyedrop), subcutaneous, intramuscular or transdermal (e.g., patch) form, all  
35 using forms well known to those of ordinary skill in the pharmaceutical

arts. An effective but non-toxic amount of the compound desired can be employed as an  $\alpha\text{v}\beta 3$  antagonist.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including  
5 type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician, veterinarian or clinician can readily determine and prescribe the effective amount of  
10 the drug required to prevent, counter or arrest the progress of the condition.

Oral dosages of the present invention, when used for the indicated effects; will range between about 0.01 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 to 10  
15 mg/kg/day, and most preferably 0.1 to 5.0 mg/kg/day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A medicament  
20 typically contains from about 0.01 mg to about 500 mg of the active ingredient, preferably, from about 1 mg to about 100 mg of active ingredient. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, compounds of the present invention may be  
25 administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal  
30 skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds  
35 herein described in detail can form the active ingredient, and are

typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as 'carrier' materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamide-phenol, or polyethyleneoxide-polylysine substituted

with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and crosslinked or amphipathic block copolymers of hydrogels.

In the schemes and examples below, various reagent symbols and abbreviations have the following meanings:

AcOH:	Acetic acid.
BH <sub>3</sub> •DMS:	Borane•dimethylsulfide.
BOC(Boc):	t-Butyloxycarbonyl.
BOP:	Benzotriazol-1-yloxytris(dimethylamino)-
	phosphonium hexafluorophosphate.
CBZ(Cbz):	Carbobenzyloxy or benzyloxycarbonyl.
CDI:	Carbonyldiimidazole.
CH <sub>2</sub> Cl <sub>2</sub> :	Methylene chloride.
CH <sub>3</sub> CN:	Acetonitrile
CHCl <sub>3</sub> :	Chloroform.
DBA:	Bis(dibenzylidene)acetone.
DEAD:	Diethyl azodicarboxylate.
DIAD:	Diisopropyl azodicarboxylate.
DIBAH or	
DIBAL-H:	Diisobutylaluminum hydride.
DIPEA:	Diisopropylethylamine.
DMAP:	4-Dimethylaminopyridine.
DME:	1,2-Dimethoxyethane.
DMF:	Dimethylformamide.
DMSO:	Dimethylsulfoxide.
DPPF:	1,1'-bis(diphenylphosphino)ferrocene.
DPFN:	3,5-Dimethyl-1-pyrazolylformamidinium nitrate.
EDC:	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide •HCl
EtOAc:	Ethyl acetate.
EtOH:	Ethanol.



	HOAc:	Acetic acid.
	HOAT:	1-Hydroxy-7-azabenzotriazole
	HOBt:	1-Hydroxybenzotriazole.
	IBCF:	Isobutylchloroformate
5	LDA:	Lithium diisopropylamide.
	MeOH:	Methanol.
	MMNG	1,1-methyl-3-nitro-1-nitrosoguanidine
	NEt <sub>3</sub> :	Triethylamine.
	NMM:	N-methylmorpholine.
10	PCA•HCl:	Pyrazole carboxamide hydrochloride.
	Pd/C:	Palladium on activated carbon catalyst.
	Ph:	Phenyl.
	pTSA	p-Toluenesulfonic acid.
	TEA:	Triethylamine.
15	TFA:	Trifluoroacetic acid.
	THF:	Tetrahydrofuran.
	TLC:	Thin Layer Chromatography.
	TMEDA:	N,N,N',N'-Tetramethylethylenediamine.
	TMS:	Trimethylsilyl.

20           The novel compounds of the present invention can be prepared according to the procedure of the following schemes and examples, using appropriate materials and are further exemplified by the following specific examples. The compounds illustrated in the examples are not, however, to be construed as forming the only genus  
25           that is considered as the invention. The following examples further illustrate details for the preparation of the compounds of the present invention. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures  
30           are degrees Celsius unless otherwise noted.

          The following Schemes and Examples describe procedures for making representative compounds of the present invention. Moreover, by utilizing the procedures described in detail in PCT International Application Publication Nos. WO95/32710, published 7  
35           December 1995, and WO95/17397, published 29 June 1995, both of which


are incorporated by reference herein in their entirety, in conjunction with the disclosure contained herein, one of ordinary skill in the art can readily prepare additional compounds of the present invention claimed herein. Additionally, for a general review describing the synthesis of

5  $\beta$ -alanines which can be utilized as the C-terminus of the compounds of the present invention, see Cole, D.C., *Recent Stereoselective Synthetic Approaches to  $\beta$ -Amino Acids*, *Tetrahedron*, 1994, 50, 9517-9582; Juaristi, E, *et al.*, *Enantioselective Synthesis of  $\beta$ -Amino Acids*, *Aldrichimica Acta*, 1994, 27, 3. In particular, synthesis of the 3-methyl- $\beta$ -alanine is

10 taught in Duggan, M.F. *et al.*, *J. Med. Chem.*, 1995, 38, 3332-3341; the 3-ethynyl- $\beta$ -alanine is taught in Zablocki, J.A., *et al.*, *J. Med. Chem.*, 1995, 38, 2378-2394; the 3-(pyridin-3-yl)- $\beta$ -alanine is taught in Rico, J.G. *et al.*, *J. Org. Chem.*, 1993, 58, 7948-7951; and the 2-amino- and 2-tosylamino- $\beta$ -alanines are taught in Xue, C-B, *et al.*, *Biorg. Med. Chem. Letts.*, 1996,

15 6, 339-344. The references described in this paragraph are all also incorporated by reference herein in their entirety.

## SCHEME 1



1-Bromo-3-(2,2-diethoxy-ethoxy)-benzene (1-2)

To a suspension of NaH (2.77 g, 115.6 mmol) in DMF (100 mL) at 0°C was added a solution of 3-bromophenol 1-1 in DMF (40 mL) over 40 min. After the addition was complete, the solution was stirred  
5 for an additional 30 min. The solution was then treated with neat bromoacetaldehyde diethyl acetal (17.36 g, 115.6 mmol). The solution was heated at 100°C for 8 h, cooled to room temperature, and extracted with Et<sub>2</sub>O (3 x 200 mL). The combined organic extracts were washed with 10% aq. NaOH (100 mL) and brine (100 mL), dried over MgSO<sub>4</sub>,  
10 filtered and concentrated to give 1-2 as a yellow oil.

TLC R<sub>f</sub> = 0.4 (10% ethyl acetate/hexanes).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.19-7.05 (m, 3H), 6.85 (d, 1H), 4.81 (t, 1H, J=6.8 Hz), 3.99 (d, 2H, J=6.8 Hz), 3.71 (m, 4H), 1.22 (t, 6H, J=7.1 Hz)

6-Bromo-benzofuran (1-3)

To a solution of the acetal 1-2 in toluene (200 mL) was added polyphosphoric acid (20 g). The biphasic mixture was heated to 100°C and stirred at this temperature for 4 h. The mixture was cooled to room temperature, poured onto ice, and extracted with Et<sub>2</sub>O (2 x 200 mL). The  
20 combined organic extracts were washed with saturated aq. NaHCO<sub>3</sub> and brine. The solution was dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (100% hexanes) to give the product 1-3 as a yellow oil.

TLC R<sub>f</sub> = 0.3 (100% hexanes).

25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.68 (s, 1H), 7.60 (d, 1H, J=2.1 Hz), 7.46 (d, 1H, J=8.4 Hz), 7.36 (dd, 1H, J=8.1, 1.5 Hz), 6.75 (dd, 1H, J=7.1, 0.9 Hz).

3-Benzofuran-6-yl-acrylic acid ethyl ester (1-4)

A mixture of the 6-bromo-benzofuran 1-3 (1.74 g, 8.79 mmol), ethyl acrylate (1.09 g, 10.98 mmol), Pd(OAc)<sub>2</sub> (0.099 g, 0.44 mmol), tri-*o*-tolylphosphine (0.268 g, 0.880 mmol), and sodium acetate (3.60 g, 43.9 mmol) in DMF (10 mL) was heated to 100°C in a sealed tube for 4 h. The mixture was cooled to room temperature, diluted with water, and extracted with Et<sub>2</sub>O (2 x 40 mL). The combined organic  
35 extracts were washed with brine (30 mL), dried over MgSO<sub>4</sub>, filtered,

and concentrated. The residue was purified by flash chromatography (10% ethyl acetate/hexanes) to give the ester 1-4 as an off-white solid.

TLC Rf = 0.3 (10% ethyl acetate/hexanes).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.78 (d, 1H, J=15.9 Hz), 7.68 (d, 1H, J=2.4 Hz), 7.66 (s, 1H), 7.59 (d, 1H, J=8.4 Hz), 7.43 (dd, 1H, J=9.0, 1.5 Hz), 6.78 (m, 1H), 6.47 (d, 1H, J=15.9 Hz), 4.27 (q, 2H, J=7.2 Hz), 1.34 (t, 3H, J=7.2 Hz).

10 3(S)-Benzofuran-6-yl-3-[benzyl-(1(R)-phenethyl)-amino]-propionic acid ethyl ester (1-5)

A solution of N-benzyl-α-(R)-methylbenzylamine (1.32 g, 6.30 mmol) in THF (25 mL) at 0°C was treated with n-BuLi (2.52 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0°C for 30 min and then cooled to -78°C. A solution of acrylate 1-4 (0.681 g, 3.15 mmol) in THF (5 mL) was added. After stirring for 15 min at -78°C, satd. aq. NH<sub>4</sub>Cl soln (5 mL) was added and the cold bath removed. The mixture was warmed to room temperature and extracted with Et<sub>2</sub>O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (10% ethyl acetate/hexanes) to give the β-aminoester 1-5 as a yellow oil.

TLC Rf = 0.8 (10% ethanol/dichloromethane).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.58 (m, 3H), 7.41 (m, 2H), 7.22 (m, 9H), 7.59 (s, 1H), 4.58 (m, 1H), 4.05 (m, 1H), 3.91 (q, 2H, J=7.1 Hz), 3.72 (m, 2H), 2.62 (m, 2H), 1.21 (d, 3H, J=7.2 Hz), 1.03 (t, 3H, J=7.1 Hz).

3(S)-Amino-3-(2,3-dihydro-benzofuran-6-yl)-propionic acid ethyl ester (1-6)


A mixture of the dibenzylamine 1-5 (1.19 g, 2.78 mmol) in EtOH/H<sub>2</sub>O/AcOH (26 mL/3 mL/1.0 mL) was degassed with argon and treated with Pd(OH)<sub>2</sub> (1.19 g). The mixture was placed under 1 atm of H<sub>2</sub>. After stirring for 18 h, the mixture was diluted with EtOAc, and filtered through celite. The filtrate was concentrated and the residue purified by flash chromatography (10% ethyl acetate/dichloromethane) to give the ester 1-6 as a white solid.


TLC R<sub>f</sub> = 0.25 (10% ethanol/dichloromethane).


<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) as the trifluoroacetate salt: δ 7.25 (d, 1H, J=8.1 Hz), 6.88 (m, 1H), 7.66 (s, 1H), 6.82 (s, 1H), 4.58 (m, 3H), 4.12 (m, 2H), 3.30 (m, 1H), 3.19 (m, 2H), 2.98 (m, 2H), 1.11 (t, 3H, J=7.2 Hz).

5

### SCHEME 2







**4-Oxo-pentanoic acid methoxy-methyl-amide (2-1)**

To a stirred solution of levulinic acid (30 g, 0.258 mol) in  
 5 CHCl3 (850 mL) at 0°C was added triethylamine (43.2 mL, 0.310 mol),  
 followed by isobutyl chloroformate (37 mL, 0.284 mol) over 15 minutes.  
 After 30 minutes, triethylamine (57.6 mL, 0.413 mol) was added, followed  
 by N,O-dimethylhydroxylamine hydrochloride (37.8 g, 0.387 mol) in 5  
 portions over 5 minutes. Vigorous bubbling ensued, and the mixture  
 10 was allowed to warm to RT and stirred for 1 h. The mixture was  
 reduced to a moist solid by rotary evaporation under reduced pressure,



slurried in 500 mL EtOAc, washed with 10% K<sub>2</sub>CO<sub>3</sub>, brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporative removal of the solvent gave 2-1 as a yellow oil. TLC R<sub>f</sub> = 0.42 (silica, 1:1 chloroform/ethyl acetate).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.74 (s, 3H), 3.18 (s, 3H), 2.65-2.95 (m, 4H),  
5 2.21 (s, 3H).

N-methoxy-N-methyl-3-(2-methyl-[1,3]dioxolan-2-yl)-propionamide (2-2)

To a solution of 2-1 (38 g, 0.239 mol) in 500 mL benzene was added ethylene glycol (17.3 mL, 0.310 mol) and p-toluenesulfonic acid (1  
10 g). The mixture was heated at reflux for 2 h with azeotropic removal of water. After cooling, the solution was washed with 200 mL sat. NaHCO<sub>3</sub>, brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporative removal of the solvent gave 2-2 as a yellow oil.

TLC R<sub>f</sub> = 0.62 (silica, ethyl acetate).

15 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.95 (m, 4H), 3.68 (s, 3H), 3.17 (s, 3H), 2.51 (t, 2H, J=8 Hz), 2.00 (t, 3H, J=6 Hz) 1.33 (s, 3H).

3-(2-Methyl-[1,3]dioxolan-2-yl)-propionaldehyde (2-3)

To a solution of 2-2 (44.74 g, 0.22 mol) in 400 mL THF at  
20 -78°C was added DIBAL (264 mL 1 M in hexanes, 0.264 mol) over 10 minutes. After stirring for 1 h, 350 ml of 1.0 M Rochelle's salt and 300 ml ether were added followed by the removal of the cooling bath. After stirring for 1 h, the organic portion was separated and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporative removal of the solvent gave 2-3 as a colorless oil.

25 TLC R<sub>f</sub> = 0.80 (silica, ethyl acetate).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 9.73 (s, 1H), 3.50 (d, 1H, J=16 Hz), 2.61 (d, 1H, J=21 Hz), 2.48 (m, 1H), 2.07 (t, 1H, J= 7H), 1.33 (s, 3H).

[3-(2-Methyl-[1,3]dioxolan-2-yl)-propylamino]-acetic acid ethyl ester (2-4)

30 To a solution of 2-3 (31.7 g, 0.22 mol) in 1000 mL 1,2-dichloroethane at 0°C were added glycine ethyl ester hydrochloride (61.5 g, 0.44 mol), triethylamine (107 mL, 0.77 mol), and NaB(OAc)<sub>3</sub>H (65.3 g, 0.308 mol). The mixture was allowed to warm to RT and stirred for 15 h. The mixture was evaporated to one-third its initial volume, diluted with

EtOAc and then washed with 10% K<sub>2</sub>CO<sub>3</sub>, brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 1:1 chloroform/ethyl acetate followed by 5% MeOH/ethyl acetate) to give 2-4 as a yellow oil.

5 TLC R<sub>f</sub> = 0.40 (silica, ethyl acetate).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 6.01 (br s, 1H), 4.21 (m, 3H), 4.03 (d, 1H, J=5 Hz), 3.93 (m, 4H), 2.62 (t, 2H, J=8 Hz), 1.53-1.67 (m, 4H), 1.29 (m, 6H).

10 [Tert-butoxycarbonyl-[3-(2-methyl-[1,3]dioxolan-2-yl)-propyl]-aminol]-acetic acid ethyl ester (2-5)

To a solution of 2-4 (24 g, 0.104 mol) in 100 mL THF were added a trace of DMAP, 20 drops of triethylamine, and BOC<sub>2</sub>O (23.8 g, 0.109 mol). After 4 h, evaporative removal of the solvent gave 2-5 as a colorless oil.

15 TLC R<sub>f</sub> = 0.38 (silica, 30% ethyl acetate/hexane).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 4.22 (m, 3H), 3.93 (m, 4H), 3.27 (m, 2H), 1.63 (m, 4H), 1.51 (s, 3H), 1.47 (s, 3H), 1.42 (s, 3H), 1.31 (s, 3H), 1.28 (m, 4H).

20 [Tert-butoxycarbonyl-(4-oxo-pentyl)-aminol]-acetic acid ethyl ester (2-6)

To a solution of 2-5 (35 g, 0.1 mol) in 600 mL acetone was added p-toluenesulfonic acid (1 g). The mixture was heated at reflux for 2 h. After cooling, the mixture was evaporated to one-fifth its initial volume, diluted with EtOAc and then washed with sat. NaHCO<sub>3</sub>, brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporative removal of the solvent gave 2-6 as a yellow oil.

25 TLC R<sub>f</sub> = 0.31 (silica, 30% ethyl acetate/hexane).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, mixture of rotamers) δ 4.20 (m, 2H), 3.92 (s, 0.85H), 3.83 (s, 1.15 H), 3.3 (m, 2H), 2.52 (m, 2H), 2.14 (s, 3H), 1.78 (m, 2H), 1.51-1.42 (3s, 9H), 1.28 (m, 3H).

30

[Tert-butoxycarbonyl-(3-[1,8]naphthyridin-2-yl)-propyl)-aminol]-acetic acid ethyl ester (2-7)

A solution of 2-6 (28 g, 97.4 mmol), 2-amino-3 formylpyridine (15.5 g, 127 mmol), proline (11.2 g, 97.4 mmol) in ethanol (250 mL) was heated at reflux for 15 h. After cooling and evaporation, the residue was chromatographed (silica gel, 1:1 chloroform/ethyl acetate) to give 2-7 as a yellow oil.

TLC  $R_f$  = 0.41 (silica, 70:25:5 chloroform/ethyl acetate/methanol)

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  9.09 (m, 1H), 8.14 (m, 2H), 7.43 (m, 2H), 4.17 (q, 2H, 7 Hz), 3.9 (2s, 2H), 3.43 (q, 2H,  $J=7$  Hz), 3.07 (m, 2H), 2.18 (m, 2H), 1.42 (s, 9H), 1.25 (m, 3H).

[(Tert-butoxycarbonyl-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino)-acetic acid ethyl ester (2-8)]

A solution of 2-7 (24.3 g, 65.1 mmol), platinum oxide (4 g) and ethanol (130 mL) was stirred under a balloon of hydrogen gas for 6 h. Following filtration and evaporation, the residue was chromatographed (silica gel, ethyl acetate) to give 2-8 as a yellow oil.

TLC  $R_f$  = 0.35 (silica, 70:25:5 chloroform/ethyl acetate/methanol)

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.05 (d, 1H,  $J=6$  Hz), 6.37 (m, 1H), 4.74 (brs, 1H), 4.18 (q, 2H,  $J=7$  Hz), 3.9 (2s, 2H), 3.32 (m, 4H), 2.63 (m, 2H), 2.51 (m, 2H), 2.72 (m, 4H), 1.43 (m, 9H), 1.26 (m, 3H).

[(Methoxy-methyl-carbamoyl)-methyl]-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-carbamic acid tert-butyl ester (2-9)]

To a solution of 2-8 (1.49 g, 3.96 mmol) in ethanol (8 mL) was added NaOH (4.36 mL 1M solution in water, 4.36 mmol). After stirring for 1 h at 50°C, HCl (4.75 mL of a 1M solution in water, 4.75 mmol) was added, and the mixture evaporated to give an oily residue. The residue was evaporated from ethanol three times, and then from acetonitrile three times, producing a yellow crusty solid which was dried under a vacuum of <2 mm Hg for 2 h. This residue was then slurried in chloroform (15 mL), and triethylamine (2.75 mL, 19.8 mmol), N,O-dimethylhydroxylamine hydrochloride (0.772 g, 7.92 mmol), HOBT (1g) and EDC (0.91 g, 4.75 mmol) were added. After stirring for 15 h, the mixture was evaporated to dryness, the residue slurried in EtOAc,

washed with sat. NaHCO<sub>3</sub>, brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporative removal of the solvent gave 2-9 as a yellow oil.

TLC R<sub>f</sub> = 0.49 (silica, 70:25:5 chloroform /ethyl acetate /methanol)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.05 (m, 1H), 6.38 (m, 1H), 4.81 (br s, 1H),

5 3.69, m, 3H), 3.37 (m, 4H), 3.18 (s, 3H), 2.64 (m, 2H), 2.53 (m, 2H), 1.88 (m, 4H), 1.44 (m, 9H).

{Tert-butoxycarbonyl-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino}-acetaldehyde (2-9A)

10 To a stirred solution of 2-9 (11.0 g, 28.0 mmol) and THF (300 ml) at -78°C was added DIBAL (1.0M/hexanes, 42 ml, 42 mmol) dropwise over 20 minutes. After 1.0 hour, 300 ml of 1.0 M Rochelle's salt was added followed by the removal of the cooling bath. The mixture was stirred for 1.0 hour and then diluted with Et<sub>2</sub>O. After 30 minutes of  
15 stirring, the organic portion was separated and dried over MgSO<sub>4</sub>. Evaporative removal of the solvent gave crude aldehyde 2-9A as a colorless oil.

TLC R<sub>f</sub> = 0.34 (silica, 75:25:5 chloroform /EtOAc/MeOH).

20 3(S)-(2-{Tert-butoxycarbonyl-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino}-ethylamino)-3-(2,3-dihydro-benzofuran-6-yl)-propionic acid ethyl ester (2-10)

A mixture of the crude aldehyde 2-9A (9.1 mmol), 1-6 (3.2 g, 11.8 mmol), powdered molecular sieves (3 g) and DCE (100 mL) was  
25 stirred for 30 minutes. The mixture was cooled to 0°C and then NaB(OAc)<sub>3</sub>H (2.7 g, 12.7 mmol) was added. After 1 hour, the reaction was diluted with EtOAc and then washed with 10% K<sub>2</sub>CO<sub>3</sub>, brine, and dried over MgSO<sub>4</sub>. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 1-3% [10:10:1 EtOH/ NH<sub>4</sub>OH/  
30 H<sub>2</sub>O]/50:50 chloroform /ethyl acetate) to give 2-10 as a yellow oil.

TLC R<sub>f</sub> = 0.23 (silica, 5% [10:10:1 EtOH/ NH<sub>4</sub>OH/ H<sub>2</sub>O]/50:50 chloroform /ethyl acetate)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.10 (d, J=7.6 Hz, 1H), 7.04 (d, J=7.3 Hz, 1H), 6.75 (m, 2H), 6.31 (d, J=7.3 Hz, 1H), 4.76 (s, 1H), 4.55 (t, J=8.6 Hz, 2H), 4.08  
35 (m, 2H), 4.00 (t, J=6.1 Hz, 1H), 3.41 (m, 2H), 3.16 (m, 6H), 2.68 (t, J=6.4 Hz,

1H), 2.59 (m, 3H), 2.48 (t, J=7.6 Hz, 2H), 1.81 (m, 4H), 1.39 (s, 9H), 1.21 (m, 3H).

5 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propylamino]-ethylamino}-propionic acid ethyl ester (2-11)

HCl gas was rapidly bubbled through a solution of 2-10 (4.0 g, 7.2 mmol) in dioxane (160 ml) at 0°C for 10 minutes. After 30 minutes, the solution was purged with argon for 30 minutes. The solution was  
10 concentrated to give the amine 2-11 as a yellow solid.  
1H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.91 (s, 1H), 7.40 (d, J=7.0 Hz, 1H), 7.23 (d, J=7.6 Hz, 1H), 7.10 (m, 2H), 6.56 (d, J=6.1 Hz, 1H), 4.58 (m, 2H), 4.04 (m, 2H), 3.49 (m, 4H), 3.19 (m, 4H), 2.90 (m, 2H), 2.79 (m, 2H), 2.30 (m, 2H), 1.98 (m, 2H), 1.85 (m, 5H), 1.15 (t, J=7.1 Hz, 3H).

15

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl ester (2-12)

To a stirred mixture of 2-11 (11.8 mmol), CH<sub>2</sub>Cl<sub>2</sub> (3 mL) and  
20 20% K<sub>2</sub>CO<sub>3</sub> was added phosgene (1.93 M toluene, 6.7 ml, 13.0 mmol) dropwise over 20 minutes. After stirring for 30 minutes, the organic layer was separated and dried over MgSO<sub>4</sub>. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 5-10% methanol /ethyl acetate) to give 2-12 as a yellow oil.  
25 TLC R<sub>f</sub> = 0.25 (silica, 70:20:10 chloroform /ethyl acetate /methanol)  
1H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.12 (d, J=7.6 Hz, 1H), 7.02 (d, J=7.3 Hz, 1H), 6.80 (d, J=7.6 Hz, 1H), 6.70 (s, 1H), 6.34 (d, J=7.3 Hz, 1H), 5.46 (t, J=7.9 Hz, 1H), 4.74 (s, 1H), 4.55 (t, J=8.9 Hz, 2H), 4.10 (q, J=7.3 Hz, 2H), 3.41 (m, 2H), 3.21 (m, 6H), 2.95 (m, 3H), 2.67 (t, J=6.1 Hz, 2H), 2.52 (t, J=7.6 Hz,  
30 2H), 1.88 (m, 5H), 1.20 (t, J=7.3 Hz, 3H).




3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (2-13)


To a solution of 2-12 (2.9g, 6.06 mmol) in EtOH (15 mL) was  
35 added 1N NaOH (7.2 ml, 7.2 mmol). After stirring for 2 h, the solvents

were evaporated and the residue chromatographed (silica gel, 25:10:1:1 followed by 15:10:1:1 ethyl acetate /EtOH /water /NH<sub>4</sub>OH) to give 2-13 as a white solid.

TLC R<sub>f</sub> = 0.24 (15:10:1:1 ethyl acetate/EtOH/water/NH<sub>4</sub>OH).

- 5 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.55 (d, J=7.3 Hz, 1H), 7.18 (d, J=7.6 Hz, 1H), 6.81 (d, J=6.1 Hz, 1H), 6.72 (s, 1H), 6.62 (d, J=7.3 Hz, 1H), 5.38 (t, J=7.9 Hz, 1H), 4.53 (t, J=8.9 Hz, 2H), 3.14 → 3.53 (9H), 2.97 (m, 3H), 2.80 (t, J=6.1 Hz, 2H), 2.67 (t, J=7.3 Hz, 2H), 1.93 (m, 4H).

SCHEME 3

SCHEME 3 (CONTINUED)

5

Ethyl 3-fluorocinnamate (3-2)

To a solution of 3-fluorobenzaldehyde 3-1 (18.16 g, 146 mmol) in dichloromethane (500 mL) was added ethyl  
 10 (triphenylphosphoranylidene)acetate (61.2 g; 176 mmol) and the resulting solution was stirred at room temperature for 18 hr. After evaporation of the solvent, the residue was swirled with ether/hexane and filtered. The filtrate was concentrated and then purified on a plug of silica gel eluting with hexane/EtOAc 9:1. Removal of the solvent  
 15 afforded the title compound 3-2 as an oil (~95% trans) which was used without further purification in the next step.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.36 (3H, t), 4.28 (2H, q), 6.43 (1H, d), 7.08 (1H, m), 7.2-7.4 (3H, m), 7.64 (1H, d).

20 N-Benzyl-(R)-α-methylbenzyl-3(S)-fluorophenyl-β-alanine ethyl ester (3-3)

To a solution of N-benzyl-(R)-α-methylbenzylamine (33.4 g, 158 mmol) in THF (450 mL) at 0°C was added n-butyllithium (1.6M in hexanes; 99 mL, 158 mmol). The dark violet solution was stirred at 0°C for 30 minutes, cooled to -78°C, and the ester 3-2 (29.2 g, 150 mmol) in  
 25 THF (100 mL) was added over 5 minutes. The resulting solution was stirred at -78°C for 1 hr., then warmed to room temperature. After 2 hrs, the mixture was poured into water and extracted with EtOAc, washed with water, then brine, dried and concentrated *in vacuo* to give



an oil. Column chromatography (silica gel; hexane/EtOAc 1:1 then pure EtOAc) gave the title compound 3-3.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.06 (3H, t), 1.28 (3H, d), 2.52 (1H, dd), 2.62 (1H, dd), 3.66 (1H, d), 3.72 (1H, d), 3.95 (2H, q), 4.44 (1H, dd), 6.95 (1H, m),  
5 7.1-7.5 (13H, m).

3(S)-Fluorophenyl-β-alanine ethyl ester hydrochloride (3-4)

A solution of the N-benzyl-(R)-α-methylbenzylamine 3-3 (28.2 g, 69.6 mmol) in ethanol (300 mL), acetic acid (30 mL) and water (3  
10 mL) was degassed with argon for 30 minutes. Pd(OH)<sub>2</sub> on carbon (20% dry weight; 2.6 g) was added and the mixture then stirred under a hydrogen atmosphere (balloon) for 2 hours. The mixture was filtered through celite and the solvent removed *in vacuo* to give an oil. This oil was dissolved in 200 mL ether, and to this solution was added 60 mL 1N  
15 HCl in ether to yield a precipitate. Filtration and washing the solid with ether/hexane gave the title compound 3-4 as a white solid.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 1.21 (3H, t), 3.0-3.2 (2H, m), 4.16 (2H, q), 4.76 (1H, t), 7.2-7.35 (3H, m), 7.5 (1H, m).


20 3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (3-5)


Compound 3-5 was prepared from 3-4 using the procedure for the preparation of 2-13.

TLC R<sub>f</sub> = 0.36 (15:10:1:1 ethyl acetate/EtOH/water/NH<sub>4</sub>OH).

25 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.41 (m, 2H), 7.09 (m, 3H), 6.54 (d, 1H, J=8.2 Hz), 5.48 (m, 1H), 3.51 (m, 2H), 3.46 (m, 3H), 3.23 (m, 2H), 2.94 (m, 2H), 2.81 (m, 4H), 2.63 (m, 2H), 1.93 (m, 2H), 1.19 (m, 2H, J=5.1 Hz).

## SCHEME 4





3-Quinolin-3-yl-propionic acid (4-2).

- A solution containing quinoline-3-carboxaldehyde 4-1 (5 g, 31.8 mmol), malonic acid (3.6 g, 35.0 mmol), and ammonium acetate (5.0 g, 63.6 mmol) in anhydrous ethanol (125 mL) was heated at reflux for 12 h. After cooling to room temperature, the resulting white solid was collected by filtration and washed with cold ethanol (150 mL) and then dried under vacuum to provide 4-2 as a white solid (3.84 g, 17.8 mmol, 56%).
- <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O): δ 8.91 (d, J = 2 Hz 1H), 8.21 (d, J = 2 Hz, 1H), 8.12 (d, J = 8 Hz, 1H), 7.84 (d, J = 7 Hz, 1H), 7.72 (t, J = 7 Hz, 1H), 7.54 (t, J = 7 Hz, 1H), 4.72 (m, 1H), 2.73 (m, 2H).

3-Phenylacetyl-amino-3-quinolin-3-yl-propionic (4-3)

- A 0°C solution of 4-2 (3.5 g, 16.2 mmol) and NaHCO<sub>3</sub> (2.7 g, 32.4 mmol) in 50% aqueous dioxane (100 mL) was treated dropwise with a solution of phenylacetyl chloride (3.00 g, 19.4 mmol) in 25 mL of dioxane. The resulting solution was stirred at 0°C for 2.5h., then warmed to room temperature, diluted with H<sub>2</sub>O (50 mL) and washed with ether (2 x 100 mL). The aqueous layer was adjusted to pH = 3 with 3N HCl and then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 150 mL). The pooled organic extracts were dried, filtered and concentrated to afford 4-3 as an off-white solid.
- <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD): δ 8.85 (d, J = 2 Hz 1H), 8.20 (d, J = 2 Hz, 1H), 8.00 (d, J = 8 Hz, 1H), 7.86 (d, J = 7 Hz, 1H), 7.76 (t, J = 7 Hz, 1H), 7.52 (t, J

= 7 Hz, 1H), 7.28 (m, 6H), 5.53 (t, J = 6.8 Hz, 1H), 3.57 (s, 2H), 2.96 (m, 2H).

3-(S)-Quinolin-3-yl-propionic acid dihydrochloride (4-6)

5                    Acid 4-3 (5.0 g, 15 mmol) was suspended in water (3.5 L), then treated with 1N NaOH (15 mL) to afford a clear solution. Penicillin amidase (Sigma, EC 3.5.1.11, 10,000 U) in 0.1 M phosphate buffer was added. The pH of the mixture was adjusted to 7.8 with 1N NaOH, and the solution was stirred at room temperature for 4 days. The reaction  
10                   was monitored periodically by HPLC and the reaction stopped once the 50% conversion was reached. Next, the reaction solution was cooled to 0°C and adjusted to pH = 3 with 3N HCl. An oily yellow precipitate formed and was collected by filtration then washed with water to afford crude 4-5 (1.8 g, 5.3 mmol). The filtrate was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x  
15                   500 mL) to afford additional 4-5 contaminated by phenylacetic acid. Both batches of crude 4-5 were combined and stirred in 3 N HCl (200 mL) at 50° for 12 h., then cooled, washed with ether (2 x 100 mL) and evaporated to afford 4-6.

20                   3-(S)-Quinolin-3-yl-propionic acid ethyl ester dihydrochloride (4-7).


                    The resolved acid 4-6 was converted to 4-7 by refluxing in ethanolic HCl.

<sup>1</sup>H NMR (300 MHz CD<sub>3</sub>OD): δ 9.25 (d, J = 2 Hz 1H), 8.31 (d, J = 2 Hz, 1H), 8.15 (d, J = 8 Hz, 1H), 7.84 (d, J = 7 Hz, 1H), 7.72 (t, J = 7 Hz, 1H), 7.54 (t, J  
25                   = 7 Hz, 1H), 4.72 (m, 1H), 4.15 (q, J = 6 Hz, 2H), 2.73 (m, 2H) 1.18 (t, J = 6 Hz, 3H).

3(S)-(Quinolin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (4-8).

30                   Compound 4-8 was prepared from 4-7 using the procedure for the preparation of 2-13.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.83 (m, 1H), 8.32 (m, 1H), 8.02 (m, 2H), 7.78 (m, 1H), 7.63 (m, 1H), 7.43 (d, 1H, J=7.3 Hz), 6.57 (d, 1H, J=7.3 Hz), 5.76 (m, 1H), 3.73 (q, 1H, J=8.2 Hz), 3.48 (m, 3H), 3.32 (m, 4H), 3.17 (m,  
35                   2H), 2.95 (m, 1H), 2.84-2.62 (, 6H), 2.10 (m, 1H), 1.88 (m, 3H).

**SCHEME 5**


**5** 3(S)-(Pyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (5-2)

Compound **5-2** was prepared from **5-1** (for preparation, see Zablocki et al., *J. Med. Chem.* 1995, 38, 2378) using the procedure for the preparation of **2-13**.




<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.54 (m, 1H), 8.47 (m, 1H), 7.85 (d, 1H, J=7.9 Hz), 7.46 (m, 2H), 6.55 (d, 1H, J=7.3 Hz), 5.57 (m, 1H), 3.63 (m, 2H), 3.46 (m, 3H), 3.18 (m, 2H), 3.01 (m, 2H), 2.77 (m, 4H), 2.60 (m, 2H), 2.05 (m, 1H), 1.93 (m, 3H).


TLC (silica): R<sub>f</sub>=0.09 (15 EtOAc/10 EtOH/1 NH<sub>4</sub>OH/ H<sub>2</sub>O)

## SCHEME 6



## SCHEME 6 (CONTINUED)




SCHEME 6 (CONTINUED)






5 The imidazolidine-ring methylated compound 6-13 was prepared using the procedure described in Scheme 2 by replacing the glycine ethyl ester with alanine ethyl ester.








In further embodiments, other imidazolidine-ring substituted systems are prepared using the procedure described in Scheme 2 by employing the desired naturally-occurring or non-naturally-occurring amino acid.



SCHEME 7

Pd(OAc)<sub>2</sub>, Et<sub>3</sub>N,  
tri-o-tolylphosphine  
ethyl acrylate, CH<sub>3</sub>CN





5-Bromo-2-methoxypyridine (7-2)

To a solution of KOH (4.2 g, 0.075 mol) in water (750 mL) was added 2-methoxypyridine 7-1 (16.4 g, 0.15 mol) followed by a dropwise addition of bromine (24 g, 0.15 mol) in 1N aqueous KBr (750 mL), and the  
5 resulting solution was stirred at room temperature for 5 hr. Solid NaHCO<sub>3</sub> was added until basic, and the solution was extracted with CHCl<sub>3</sub> (3x500 mL). The organic layer was washed with 10% NaHSO<sub>3</sub>, then brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and the solvent removed in vacuo. The resulting dark brown oil was predominantly the desired compound  
10 7-2 and was used as such in the next step.  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.91 (3H, s), 6.66 (1H, d), 7.62 (1H, dd), 8.20 (1H, d).

Ethyl 3-(6-methoxypyridin-3-yl)acrylate (7-3)

15 A solution of the 5-bromo-2-methoxypyridine 7-2 (74.3 g, 0.4 mol), ethyl acrylate (150 mL, 1.4 mol), triethylamine (150 mL, 1.08 mol), palladium acetate (10 g, 0.045 mol) and tri-*o*-tolylphosphine (20 g, 0.066 mol) in 100 mL acetonitrile was degassed with argon for 10 minutes. The mixture is heated at 90°C for 12 hr, then the volatiles were removed  
20 in vacuo. Toluene (300 mL) was added and the mixture concentrated again. Diethyl ether (300 mL) was added and the mixture filtered through a pad of silica gel eluting with 800 mL of diethyl ether. After removal of the diethyl ether, the residue was chromatographed on silica gel eluting with EtOAc/hexane 1:19 then 1:14 then 1:9 to give 7-3 as a  
25 yellow solid.  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.34 (3H, t), 3.97 (3H, s), 4.26 (2H, q), 6.34 (1H, d), 6.76 (1H, d), 7.63 (1H, d), 7.77 (1H, dd), 8.27 (1H, d).

N-Benzyl-(R)-α-methylbenzyl-3(S)-(6-methoxypyridin-3-yl)-β-alanine ethyl ester (7-4)  
30

To a solution of N-benzyl-(R)-α-methylbenzylamine (97.5 g, 462 mmol) in THF (750 mL) at 0°C was added n-butyllithium (2.5M in hexanes; 178.5 mL, 446 mmol). The dark violet solution was stirred at 0°C for 20 minutes, cooled to -78°C and the ester

7-3 (63.7 g, 308 mmol) in THF (250 mL) was added over 60 minutes. The resulting solution was stirred at -78°C for 1 hr., then cannulated into saturated NH<sub>4</sub>Cl and extracted with EtOAc, washed with water then brine, dried and concentrated *in vacuo* to give an oil. Column chromatography (silica gel; hexane/EtOAc 9:1 then 4:1) gave 7-4 as an oil contaminated with N-benzyl-(R)- $\alpha$ -methylbenzylamine. This oil was taken up in 5% AcOH in water and extracted with diethyl ether (4x). The organic layers were dried over MgSO<sub>4</sub> and the solvent removed to give the title compound 7-4.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  1.08 (3H, t), 1.27 (3H, d), 2.52 (1H, dd), 2.62 (1H, dd), 3.66 (1H, d), 3.70 (1H, d), 3.93 (3H, s), 3.95 (2H, m), 4.41 (1H, dd), 6.74 (1H, d), 7.15-7.45 (10H, m), 7.64 (1H, dd), 8.15 (1H, d).

3(S)-(6-methoxypyridin-3-yl)- $\beta$ -alanine ethyl ester (7-5)

To a degassed (argon) solution of the ester 7-4 (70 g) in EtOH (250 mL), HOAc (25 mL) and water (2 mL) was added 20% Pd(OH)<sub>2</sub> on carbon. The mixture was placed under hydrogen using a balloon and the resulting mixture was stirred for 24 hr. After filtration through celite (washing with EtOAc), the solvent was removed *in vacuo* to afford a waxy solid. This was dissolved in 200 mL water and extracted with diethyl ether (2x200 mL). The aqueous layer was then treated with solid K<sub>2</sub>CO<sub>3</sub> until fully saturated and extracted with EtOAc (4x200 mL). After drying over MgSO<sub>4</sub>, the solvent was removed *in vacuo* to give the title compound 7-5 as an oil which solidified in the freezer.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  1.23 (3H, t), 2.61 (1H, dd), 2.68 (1H, dd), 3.92 (3H, s), 4.15 (2H, q), 4.41 (1H, dd), 6.93 (1H, d), 7.62 (1H, dd), 8.13 (1H, d).

3(S)-(2-(tert-Butoxycarbonyl)-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino)-ethylamino)-3-(2-methoxypyridin-5-yl)-propionic acid ethyl ester (7-6)

To a solution of the amine 7-5 (6.85 g, 30.5 mmol) in 2-propanol (300 mL) at room temperature was added acetic acid (1.75 mL, 30.5 mmol), NaOAc (24.6 g, 0.3 mol) and 4Å molecular sieves (5 g). The aldehyde 2-9A (8.1 g, 24.3 mmol) in 2-propanol (150 mL) was added and

the mixture stirred for 15 minutes, then cooled to 0°C and NaCNBH<sub>3</sub> (5.66 g, 90 mmol) added in one lot. The resulting mixture was allowed to warm to room temperature and stirred for 16 hr before being filtered through celite. After removal of the solvent in vacuo, the residue was  
5 treated with 10% aqueous KHSO<sub>4</sub> for 30 minutes, basified with solid K<sub>2</sub>CO<sub>3</sub> (to pH ~10) and extracted with 3x 200 mL EtOAc. The EtOAc layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated *in vacuo* to give an oil. Column chromatography (silica gel; 5% MeOH in CHCl<sub>3</sub>) gave 7-6 as an oil contaminated with ~7% of the β-alanine.

10 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.20 (3H, t), 1.42 (9H, s), 1.7-2 (4H, br m), 2.5-2.8 (8H, m), 3.2 (4H, m), 3.42 (2H, m), 3.92 (3H, s), 4.06 (2H, q), 5.0-5.4 (1H, bs), 6.36 (1H, br s), 6.72 (1H, d), 7.12 (1H, br s), 7.58 (1H, dd), 8.07 (1H, d).

15 3(S)-(6-Methoxypyridin-3-yl)-3-{2-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propylamino]-ethylamino}-propionic acid ethyl ester (7-7)

A solution of the ester 7-6 (14.1 g, 26 mmol) in EtOAc (350 mL) cooled to -20°C was treated with HCl (gas) for 10 minutes then stoppered and stirred at 0°C for 1.5 hr. The volatiles were removed in  
20 vacuo, the residue taken up in 150 mL of water and treated with solid K<sub>2</sub>CO<sub>3</sub> to pH~10. This solution was extracted with EtOAc (3x 150 mL), washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give an oil. Column chromatography (silica gel; 5% MeOH in CHCl<sub>3</sub>) gave the β-alanine 7-5; further elution with 5% MeOH in CHCl<sub>3</sub> saturated with  
25 NH<sub>3</sub> gave 7-7 as a viscous oil.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.20 (3H, t), 1.8-1.95 (4H, m), 2.5-2.8 (12H, m), 3.39 (2H, m), 3.92 (3H, s), 4.09 (2H, q), 5.01 (1H, bs), 6.34 (1H, d), 6.72 (1H, d), 7.06 (1H, d), 7.59 (1H, dd), 8.07 (1H, d).

30 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl ester (7-8)

To a solution of the diamine 7-7 (8.03 g, 18.2 mmol), DIPEA (9.5 mL, 54.6 mmol) and DMAP (250 mg) in 1,2-dichloroethane (150 mL)

cooled to -20°C was added p-nitrophenyl chloroformate (3.85 g, 19.1 mmol) in 1,2-dichloroethane (25 mL) dropwise such that the internal temperature remains below -15°C. The resulting mixture was allowed to warm to 0°C and stirred for 45 minutes, then heated to reflux for 4 hr.

5 After cooling, the solvent was evaporated in vacuo, the residue taken up in EtOAc and washed successively with 10% K<sub>2</sub>CO<sub>3</sub> (6x 150 mL) and brine. The EtOAc layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated *in vacuo* to give an oil (6.27 g). Column chromatography (silica gel; 5% EtOH in CH<sub>2</sub>Cl<sub>2</sub>) gave 7-8 as an oil.

10 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.20 (3H, t), 1.8-1.95 (4H, m), 2.52 (2H, dd), 2.68 (1H, dd), 2.9-3.1 (3H, m), 3.15-3.3 (5H, m), 3.39 (2H, m), 3.92 (3H, s), 4.11 (2H, q), 4.8 (1H, bs), 5.42 (1H, t), 6.34 (1H, d), 6.72 (1H, d), 7.03 (1H, d), 7.60 (1H, dd), 8.08 (1H, d).

15 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (7-9)

To a solution of the ester 7-8 (3.48 g, 7.4 mmol) in MeOH (50 mL) and water (30 mL) at room temperature was added 1N NaOH solution (22.3 mL, 22.3 mmol) and the mixture stirred for 16 hr. After  
20 removal of the solvent in vacuo, the residue was treated with 25 mL 1N HCl and the solvent removed again. Column chromatography of the residue (silica gel; EtOAc/EtOH/aq. NH<sub>4</sub>OH/H<sub>2</sub>O 20:10:1:1 then 15:10:1:1) gave a gum which was crystallized from a minimum amount of water and filtered to give 7-9 as a white solid.


25 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 1.75-2.1 (4H, m), 2.55-3.1 (8H, m), 3.28 (1H, q), 3.3 (1H, m), 3.4-3.55 (3H, m), 3.63 (1H, q), 3.85 (3H, s), 5.47 (1H, dd), 6.55 (1H, d), 6.80 (1H, d), 7.48 (1H, d), 7.68 (1H, d), 8.09 (1H, d).

In a further embodiment, the R-enantiomer, i.e. 3(R)-(6-methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid was prepared by substituting N-benzyl-(S)-α-methylbenzylamine for the N-benzyl-(R)-α-methylbenzylamine in preparing intermediate 7-4.

30

In yet a further embodiment, the racemate, i.e. 3-(6-methoxypyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid was prepared by substituting racemic N-benzyl- $\alpha$ -methylbenzylamine for the N-benzyl-  
 5 (R)- $\alpha$ -methylbenzylamine in preparing intermediate 7-4.

### SCHEME 8



#### 5-Bromo-2-ethoxypyridine (8-2)

10 Sodium metal (4.87 g, 0.212 mol) was added to ethanol (200 mL) and stirred until completely dissolved. To this solution was added 2,5-dibromopyridine 8-1 (Aldrich; 10 g, 0.0424 mol) and the resulting mixture was stirred at reflux for 16 hr. The solvent was removed in vacuo and the residue partitioned between water and EtOAc. After  
 15 extraction with EtOAc (2x), the organic layer was washed with brine, dried (MgSO<sub>4</sub>) and concentrated to give 8-2 as a red-brown solid which was used as such in the next step.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  1.4 (3H, t), 4.33 (2H, q), 6.63 (1H, d), 7.62 (1H, dd), 8.19 (1H, d).

20

3(S)-(6-Ethoxypyridin-3-yl)- $\beta$ -alanine ethyl ester (8-3)

The title compound 8-3 was prepared from 8-2 using the procedure described for the synthesis of 7-5.


<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  1.25 (3H, t), 1.39 (3H, t), 2.61 (1H, dd), 2.67  
5 (1H, dd), 4.15 (2H, q), 4.34 (2H, q), 4.40 (1H, dd), 6.71 (1H, d), 7.62 (1H, dd),  
8.11 (1H, d).

3(S)-(6-Ethoxypyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-  
1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (8-4)

10 The title compound 8-4 was prepared from 2-9A and 8-3 using the procedure described in Scheme 2.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD)  $\delta$  1.37 (3H, t), 1.8-2.0 (4H, m), 2.65 (2H, t),  
2.82 (2H, t), 2.95-3.10 (2H, m), 3.15 (2H, m), 3.23 (2H, dt), 3.46 (2H, m), 3.51  
(2H, t), 4.32 (2H, q), 5.41 (1H, t), 6.62 (1H, d), 6.84 (1H, d), 7.57 (1H, d), 7.76  
15 (1H, dd), 8.13 (1H, d).



**SCHEME 9****3-(6-Amino-pyridin-3-yl)-acrylic acid *tert*-butyl ester (9-2)**

A mixture of 2-amino-5-bromo-pyridine (**9-1**) (10 g, 58 mmol), *tert*-butyl acrylate (50 mL, 344 mmol), triethylamine (50 mL, 359 mmol), tri-*o*-tolylphosphine (3.0 g, 9.8 mmol) and  $\text{Pd}(\text{OAc})_2$  (1.0 g, 4.5 mmol) in 150 mL  $\text{CH}_3\text{CN}$  was purged with argon for 5 min and subsequently refluxed at  $110^\circ\text{C}$  for 20 hr. The mixture was then cooled and concentrated. The residue was purified using silica gel flash

chromatography (EtOAc/hexanes 1:1) to afford the desired product 9-2 as a solid.

R<sub>f</sub> (silica, EtOAc/hexanes 1:1) = 0.26

5    3(S)-(6-Amino-pyridin-3-yl)-3-[benzyl-(1(R)-phenylethyl)-amino]-propionic acid *tert*-butyl ester (9-3)

To a cooled (0°C) solution of (R)-(+)-N-benzyl- $\alpha$ -methylbenzylamine (4.0 g, 19 mmol) in 50 mL THF was gradually added n-butyllithium (11.3 mL, 2.5 M, 28.2 mmol) over 5 min. The mixture was stirred for 30 min at 0°C and cooled to -78°C. A solution of 9-2 (2.0 g, 9.4 mmol) in 20 mL THF was gradually added. After stirring for 40 min at -78°C, it was treated with NH<sub>4</sub>Cl (sat.) at -78°C, warmed to room temperature and extracted three times with EtOAc. The combined organic layers were washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. After solvent evaporation, the residue was purified using silica gel flash chromatography (EtOAc/hexanes, 1:2) to afford the desired product 9-3 as an oil.

R<sub>f</sub> (silica, EtOAc/hexanes 1:1) = 0.28

20    3(S)-Amino-3-(6-amino-pyridin-3-yl)-propionic acid ethyl ester•2 HCl (9-4)

A mixture of 9-3 (0.5 g, 1.2 mmol) and 10% Pd/C (0.4 g) in 10 mL AcOH was purged with argon for 5 min and then heated at 78°C. 1,4-Cyclohexadiene (2 mL, 21.1 mmol) was then gradually added. The reaction mixture was stirred for 3 hr and filtered through a celite pad. The solution was concentrated and the residue was purified using silica gel flash chromatography (EtOAc/MeOH/NH<sub>4</sub>OH 1 :1:0.04) to afford an oil. To the oil (1.2 g) in 20 mL EtOH was introduced HCl gas for 10 min. The mixture was stirred 24 hr and then concentrated to afford the desired product 9-4 as the HCl salt.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD)  $\delta$  8.11 (d, J=9.6 Hz, 1H), 8.08 (s, 1H), 7.13 (d, J=9.6 Hz, 1H), 4.77 (m, 1H), 4.18 (q, J=6.8 Hz, 2H), 3.22-3.02 (m, 2H), 1.24 (t, J=6.8 Hz, 3H).


3(S)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid (9-5)


The title compound 9-5 was prepared as the TFA salt from 2-9A and 9-4 using the procedure described in Scheme 2.

- 5 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.90 (d, J=2.1 Hz, 1H), 7.46 (dd, J=8.7, 2.1 Hz, 1H), 7.03 (d, J=8.7 Hz, 1H), 6.47 (d, J=8.7 Hz, 1H), 6.34 (d, J=8.7 Hz, 1H), 5.38-5.30 (m, 1H), 3.40-3.37 (m, 2H), 3.26-3.16 (m, 4H), 3.03-2.86 (m, 2H), 2.70-2.66 (m, 2H), 2.55-2.50 (m, 2H), 2.14-2.02 (m, 2H), 1.93-1.79 (m, 4H).

10

## SCHEME 10





3-(3-Hydroxy-4-nitrophenyl)-acrylic acid ethyl ester (10-2)

To a stirred solution of aldehyde 10-1 (20.28 g, 132.5 mmol) in  
 5  $\text{CH}_2\text{Cl}_2$  (400 mL) at room temperature was added  
 (carbethoxymethylene)triphenylphosphorane (46.12 g, 132.5 mmol) over  
 a 10 min period. The resulting orange solution was stirred at room  
 temperature for 2 h. The solution was concentrated to one-fourth its  
 volume. Flash chromatography (silica gel; 30:70 EtOAc/hexanes) gave  
 10 the title compound 10-2 as a bright yellow solid.  
 TLC  $R_f = 0.75$  (25:75 EtOAc/hexanes)  
 $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.14 (d, 1H), 7.60 (d, 1H), 7.15 (dd, 1H), 6.54  
 (d, 1H), 4.30 (q, 2H), 1.36 (t, 3H).

3-(4-Amino-3-hydroxyphenyl)-acrylic acid ethyl ester (10-3)

To a stirred suspension of 10-2 (4.64 g, 19.6 mmol), NH<sub>4</sub>Cl (524 mg, 9.8 mmol), EtOH (140 mL) and H<sub>2</sub>O (70 mL) was added iron dust (2.72 g, 48.9 mmol). The resulting yellow suspension was refluxed for 1.5 h. and then the solution was filtered while hot through celite. The filtrate was concentrated and the residue was partitioned between EtOAc and brine. The layers were separated and the EtOAc layer dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give 10-3 which was used without further purification in the next step.

10 TLC R<sub>f</sub> = 0.2 (25:75 EtOAc/hexanes)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.57 (d, 1H), 7.00 (m, 2H), 6.68 (d, 1H), 6.20 (d, 1H), 4.26 (q, 2H), 4.10 (bs, 2H), 1.33 (t, 3H).

3-[4-(2-Chloroacetyl-amino)-3-hydroxyphenyl]acrylic acid ethyl ester (10-4)

To a stirred solution of 10-3 (3.38 g, 16.3 mmol) in CHCl<sub>3</sub> (80 mL) was added saturated NaHCO<sub>3</sub> (50 mL) and it was then chilled to 0°C. A solution of chloroacetyl chloride (1.94 mL, 24.4 mmol) in CHCl<sub>3</sub> (30 mL) was added dropwise to the chilled biphasic. Upon addition completion, the reaction was stirred at 0°C for 1 h. The layers were separated and the aqueous layer was extracted twice with EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give 10-4 which was used without further purification in the next step.

25 TLC R<sub>f</sub> = 0.4 (25:75 EtOAc/hexanes)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 10.33 (s, 1H), 9.58 (s, 1H), 8.02 (d, 1H), 7.51 (d, 1H), 7.19 (d, 1H), 7.12 (s, 1H), 6.39 (d, 1H), 4.42 (s, 2H), 4.17 (q, 2H), 1.25 (t, 3H).

3-(3-Oxo-3,4-dihydro-2H-benzof[1,4]oxazin-7-yl) acrylic acid ethyl ester (10-5)

To a stirred solution of 10-4 (4.28 g, 15.0 mmol) in DMF (50 mL) was added K<sub>2</sub>CO<sub>3</sub> (4.50 g, 32.6 mmol). The resulting suspension was heated to 50°C for 12 h., after which time the reaction was concentrated. The residue was partitioned between saturated NaHCO<sub>3</sub>

and EtOAc and extracted twice with EtOAc. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated. Flash chromatography (silica gel; 25:75 EtOAc/hexanes) yielded 10-5 as a beige solid.

5 TLC R<sub>f</sub> = 0.5 (25:75 EtOAc/hexanes).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.91 (s, 1H), 7.54 (d, 1H), 7.37 (s, 1H), 7.31 (d, 1H), 6.90 (d, 1H), 6.51 (d, 1H), 4.60 (s, 2H), 4.16 (q, 2H), 1.24 (t, 3H).

10 3(R)-[Benzyl-(1-phenylethyl)-aminol-3-(S)-(3-oxo-3,4-dihydro-2H-benzol[1,4]oxazin-7-yl) propionic acid ethyl ester (10-6)

To a stirred solution of (R)-(+)-N-benzyl-α-methylbenzylamine (5.43 g, 25.7 mmol) and anhydrous THF (75 mL) at 0°C was added butyllithium (10.3 mL, 2.5 M/hexanes, 25.7 mmol) via syringe. The violet-red solution was stirred at 0°C for 15 minutes and then cooled to -78°C. A solution of 10-5 (2.12 g, 8.6 mmol) in anhydrous THF (50 mL) was added via syringe, and the resulting brown solution was stirred at -78°C for 30 minutes. The brown solution was quenched with saturated NH<sub>4</sub>Cl, the mixture then warmed to room temperature and extracted twice with Et<sub>2</sub>O. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated. Flash chromatography (silica gel; 15:85 to 25:75 EtOAc/hexanes) yielded 10-6 as a white foam.

TLC R<sub>f</sub> = 0.25 (25:75 EtOAc/hexanes)

15 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.89 (s, 1H), 7.32 (m, 10H), 7.10 (m, 2H), 6.91 (d, 1H), 4.62 (s, 2H), 4.39 (m, 1H), 4.13 (q, 2H) 3.96 (m, 1H), 3.68 (s, 2H), 2.56 (m, 2H), 1.28 (m, 6H).

30 3(R)-[Benzyl-(1-phenylethyl)-aminol-3-(S)-(4-methyl-3-oxo-3,4-dihydro-2H-benzol[1,4]oxazin-7-yl) propionic acid ethyl ester (10-7)

To a stirred suspension of NaH (65 mg, 60%, 1.6 mmol) in DMF (5 mL) under argon was added a solution of 10-6 (650 mg, 1.4 mmol) in DMF (10 mL) via syringe. This yellow solution was stirred at room temperature for 30 minutes. Iodomethane (0.5 mL, 8.0 mmol) was added and the solution then stirred at room temperature for an additional 30 minutes. The reaction was quenched with saturated

NaHCO<sub>3</sub>. The aqueous layer was extracted three times with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated. Flash chromatography (silica gel; 25:75 EtOAc/hexanes) afforded 10-7 as a clear oil.

- 5    TLC R<sub>f</sub> = 0.6 (25:75 EtOAc/hexanes)  
     <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.30 (m, 10H), 7.06 (m, 2H), 6.91 (d, 1H), 4.62 (s, 2H), 4.39 (m, 1H), 4.13 (q, 2H) 3.96 (m, 1H), 3.68 (s, 2H), 3.35 (s, 3H), 2.56 (m, 2H), 1.26 (m, 6H).

- 10   3(S)-Amino-3-(4-methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)propionic acid ethyl ester (10-8)

- A stirred solution of 10-7 (581 mg, 1.2 mmol), MeOH (10 mL), AcOH (1.0 mL), and H<sub>2</sub>O (0.3 mL) was degassed with argon for 5 minutes. Pd(OH)<sub>2</sub> (581 mg) was added and the reaction was placed  
15   under 1 atm of H<sub>2</sub> for 2.5 h. The reaction was diluted with EtOAc and filtered through celite. The filtrate was concentrated to yield 10-8 as a clear oil.

- TLC R<sub>f</sub> = 0.3 (5:95 MeOH/CH<sub>2</sub>Cl<sub>2</sub>)  
     <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.04 (m, 2H), 6.93 (dd, 1H), 4.61 (s, 2H), 4.39  
20   (m, 1H), 4.13 (q, 2H), 3.37 (b, 2H), 3.35 (s, 3H), 2.69 (m, 2H), 1.24 (t, 3H).

3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (10-9)


- 25        The title compound 10-9 was prepared from 2-9A and 10-8 using the procedure described in Scheme 2.

- <sup>1</sup>H NMR (400 MHz, d<sub>6</sub>-DMSO) δ 7.83 (bs, 1H), 7.60 (d, 1H), 7.11 (d, 1H), 6.99 (d, 1H), 6.93 (s, 1H), 6.63 (d, 1H), 5.20 (t, 1H), 4.64 (s, 2H), 3.3-2.8 (m, 10H), 3.25 (s, 3H), 2.72 (m, 2H), 2.59 (m, 2H), 1.81 (m, 2H), 1.74 (m, 2H).

30



## SCHEME 11



2-tert-Butoxycarbonylamino-5-aminopyridine (11-2)

A solution of 2-amino-4-bromopyridine 11-1 (10.1 g, 58.4 mmol) in 150 mL of melted *t*-BuOH was treated with di-*tert*-butyl dicarbonate (14.0 g, 64.2 mmol). After the solution was stirred for 12 hr, the solvent was evaporated. The residue was purified using silica gel flash chromatography (CHCl<sub>3</sub>/hexanes, 5:1) to afford the desired product 11-2 as a solid.

R<sub>f</sub> (silica, 100% CHCl<sub>3</sub>) = 0.56

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.82 (bs, 1H), 8.38 (d, 1H), 8.78 (d, 1H), 7.78 (dd, 1H), 1.55 (s, 9H).

2-(tert-Butoxycarbonyl-methyl-amino)-5-aminopyridine (11-3)

To a solution of 11-2 (6.0 g, 22.0 mmol) in 50 mL DMF at 0°C was added NaH gradually. After the mixture was stirred for 40 min, CH<sub>3</sub>I (3.4 g, 24.0 mmol) was added in one portion. The reaction mixture was stirred for 5 hr, treated with 300 mL water and extracted three times with ethyl ether. The combined organic layers were washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. After solvent removal, the residue was purified by silica gel flash chromatography (CHCl<sub>3</sub>/hexanes 6:1) to afford the desired product 11-3 as a solid.

R<sub>f</sub> (silica, 100% CHCl<sub>3</sub>) = 0.40

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.40 (dd, 1H), 7.68 (m, 2H), 3.36 (s, 3H), 1.55 (s, 9H).

3-[6-(tert-Butoxycarbonyl-methyl-amino)-pyridin-3-yl]-acrylic acid ethyl ester (11-4)

A mixture of 11-3 (6.0 g, 20.9 mmol), ethyl acrylate (6.3 mL, 62.7 mmol), triethylamine (17 mL, 125.5 mmol), tri-*o*-tolylphosphine (1.3 g, 6.2 mmol) and Pd(OAc)<sub>2</sub> (0.5 g, 2.1 mmol) in 50 mL CH<sub>3</sub>CN was purged with argon for 5 min and subsequently refluxed at 110°C for 20 hr. The mixture was cooled and concentrated. The residue was purified using silica gel flash chromatography (EtOAc/hexanes 1:3) to afford the desired product 11-4 as an oil.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.47 (bs, 1H), 7.82 (m, 2H), 7.64 (d, 1H), 6.42 (d, 1H), 4.27 (q, 2H), 3.43 (s, 3H), 1.54 (s, 9H), 1.34 (t, 3H).

3-Benzylamino-3-[6-(tert-butoxycarbonyl-methyl-amino)-pyridin-3-yl]-propionic acid ethyl ester (11-5)


5 A mixture of 11-4 (1.7 g, 5.6 mmol) and benzylamine (8 mL, 73.2 mmol) was heated in a sealed-tube at 95°C for 24 hr. The crude reaction mixture was purified using silica gel flash chromatography (EtOAc/hexanes 1:3 to 1:1) to afford the desired product 11-5 as an oil. R<sub>f</sub> (silica, EtOAc/hexanes 1:1) = 0.63.

10 3-Amino-3-[6-(tert-butoxycarbonyl-methyl-amino)-pyridin-3-yl]-propionic acid ethyl ester (11-6)

A mixture of 11-5 (1.5 g 3.6 mmol), 20% Pd(OH)<sub>2</sub>/C (0.3 g), AcOH (5.5 mL) and EtOH (50 mL) was purged with argon 3 times under vacuum. The reaction mixture was stirred under balloon hydrogenation  
15 condition for 16 hr and filtered through a celite pad. After solvent removal, the desired product 11-6 was obtained as the acetate salt. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.38 (d, 1H), 7.70 (m, 2H), 4.50 (dd, 1H), 4.15 (q, 2H), 3.40 (s, 3H), 2.80 (m, 2H), 1.25 (t, 3H).

20 3-(6-Methylamino-pyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (11-7)

The title compound 11-7 was prepared as the TFA salt from 2-9A and 11-6 using the procedure described in Scheme 2.  
25 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.92 (dd, J=9.6, 1.6 Hz, 1H), 7.79 (d, J=1.6 Hz, 1H), 7.57 (d, J=7.6 Hz, 1H), 7.03 (d, J=9.6 Hz, 1H), 6.63 (d, J=7.6 Hz, 1H), 5.28 (m, 1H), 3.51-3.36 (m, 5H), 3.28-3.17 (m, 3H), 3.05 (m, 2H), 3.02 (s, 3H), 2.82 (m, 2H), 2.67 (m, 2H), 1.98-1.84 (m, 4H).

SCHEME 123-(2-Fluoro-biphenyl-4-yl)-acrylic acid ethyl ester (12-2)

A solution of 2-fluoro-4-bromobiphenyl 12-1 (7.5 gm, 31.8 mmol), ethyl acrylate (4.3 mL), Pd(OAc)<sub>2</sub> (0.714 gm, 3.2 mmol), tri-o-tolylphosphine (1.94 gm, 1.5 mmol), and triethylamine (12 mL) was

heated to 100°C in a sealed tube for 12 h. The reaction was cooled to room temperature and diluted with dichloromethane (40 mL). The organic solution was washed with 10% aq. citric acid (20 mL), satd. aq. NaHCO<sub>3</sub>, and brine (20 mL). The organic solution was dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (95:5 to 90:10 hexanes/EtOAc) to give the acrylate ester 12-2 as a white solid.

TLC R<sub>f</sub> = 0.44 (10% ethyl acetate/hexanes).

10 3-[Benzyl-(1(R)-phenylethyl)-aminol-3-(2-fluoro-biphenyl-4-yl)-propionic acid ethyl ester (12-3)

A cooled (0°C) solution of (R)-(+)-N-benzyl- $\alpha$ -methylbenzylamine (8.9 mL, 42.6 mmol) in THF (100 mL) was treated with n-butyllithium (26.6 mL of a 1.6 M soln in hexanes; 42.6 mmol). After stirring for 10 min, the purple solution was cooled to -78°C and treated with a solution of ester 12-2 (5.76 g, 21.3 mmol) in THF (10 mL). After stirring for 20 min, the solution was quenched with satd aq NH<sub>4</sub>Cl soln (5 mL), and the cold bath removed. The reaction mixture was diluted with Et<sub>2</sub>O (100 mL), and washed with 10% aq citric acid (50 mL), satd aq NaHCO<sub>3</sub> (50 mL), 5% aq acetic acid (30 mL), 10% aq K<sub>2</sub>CO<sub>3</sub> (50 mL), and brine (50 mL). The solution was dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (90:10 hexanes/EtOAc) to give adduct 12-3.

TLC R<sub>f</sub> = 0.48 (10% ethyl acetate/hexanes).

25

3-Amino-3-(2-fluoro-biphenyl-4-yl)-propionic acid ethyl ester (12-4)

A solution of the dibenzylamine 12-3 (5.65 gm, 11.75 mmol) in EtOH/HOAc (90/10 mL) was purged with argon and treated with Pd(OH)<sub>2</sub> (3 g) and placed under 1 atm of H<sub>2</sub> gas for 12 h. Additional portions (2.5 g) of Pd(OH)<sub>2</sub> were added after 24 h, 48 h and 144 h. The reaction mixture was purged with argon, filtered through Celite, and the filtrate dissolved in aq HCl (pH=1). The aqueous solution was washed with EtOAc, neutralized with satd aq NaHCO<sub>3</sub>, and extracted with EtOAc (3 x 30 mL). The combined organic solutions were washed

with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to give the desired product 12-4.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.41 (m, 8H), 4.10 (m, 1H), 4.06 (m, 2H), 2.73 (m, 2H), 1.18 (m, 3H) ppm.


5

3(S)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl) propionic acid (12-5)

The title compound 12-5 was prepared from 2-9A and 12-4 using the procedure described in Scheme 2.

10 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.49 (m, 9H), 6.64 (d, J= 7.3 Hz, 1H), 5.49 (m, 1H), 3.31 (m, 9H), 2.83 (m, 2H), 2.74 (m, 2H), 1.97 (m, 4H) ppm.

## SCHEME 13

3-(3-Hydroxy-4-nitro-phenyl)-acrylic acid ethyl ester (13-2)

To a solution of aldehyde 13-1 (15.0 g, 98.0 mmol) in  $\text{CH}_2\text{Cl}_2$   
 5 (300 mL) was slowly added carboethoxymethylenetriphenylphosphorane

(34.1 g, 98.0 mmol). The orange solution was stirred for 12 h at ambient temperature. The solution was concentrated to a paste and purified by flash chromatography (10% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>) to give 13-2 as a yellow solid.

TLC R<sub>f</sub> = 0.51 (30% ethyl acetate/hexanes).

- 5 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.08 (d, J=8.4 Hz, 1H), 7.63 (d, J=16.2 Hz, 1H), 7.35 (d, J=1.5 Hz, 1H), 7.27 (dd, J=8.4, 1.5 Hz, 1H), 6.65 (d, J=15.9 Hz, 1H), 4.25 (q, J=7.2 Hz, 2H), 1.32 (t, J=6.9 Hz, 3H) ppm.

3-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-acrylic acid ethyl ester (13-3)

- 10 To a solution of the nitrophenol 13-2 (12.0 g, 57.4 mmol) in warm (70°C) AcOH/H<sub>2</sub>O (200 mL) was added iron dust (9.61 g, 172.2 mmol). The brown heterogeneous mixture was stirred for 30 min at 70-80°C. The mixture was filtered hot through Celite, and the Celite bed washed with EtOAc (2 x 200 mL). The filtrate was cautiously neutralized  
15 with satd aq NaHCO<sub>3</sub> (3 x 100 mL). The solution was dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give an orange solid (9.6 g, 81%). A portion of this solid (4.5 g, 21.7 mmol) was dissolved in THF (150 mL) and treated with 1,1-carbonyldiimidazole (3.87 g, 23.8 mmol), and  
20 the solution was stirred at ambient temperature for 24 h. The solution was diluted with EtOAc (100 mL) and washed with 10% HCl (50 mL) and brine (50 mL). The solution was dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give 13-3 as a yellow solid.

- 25 TLC R<sub>f</sub> = 0.49 (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>).  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.77 (d, J=15.9 Hz, 1H), 7.55 (s, 1H), 7.41 (d, J=8.4 Hz, 1H), 7.09 (d, J=8.1 Hz, 1H), 6.47 (d, J=15.9 Hz, 1H), 4.22 (q, J=7.2 Hz, 2H), 1.31 (t, J=7.2 Hz, 3H) ppm.

- 30 3(S)-Amino-3-(2-oxo-2,3-dihydro-benzoxazol-6-yl)-propionic acid ethyl ester (13-4)

- A solution of (R)-(+)-N-benzyl-α-methylbenzylamine (4.08 g, 19.3 mmol) in THF (120 mL) at 0°C was treated with n-BuLi (7.72 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0°C for 30  
35 min and then cooled to -78°C. A solution of acrylate 13-3 (1.5 g, 6.43




mmol) in THF (20 mL) was added. After stirring for 15 min at -78°C, satd aq NH<sub>4</sub>Cl soln (25 mL) was added and the cold bath removed. The mixture was warmed to room temperature and extracted with Et<sub>2</sub>O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL),  
5 dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (30% ethyl acetate/hexanes) to give 2.74 g of the β-aminoester as a yellow oil. The aminoester was dissolved in EtOH/H<sub>2</sub>O/AcOH (54 mL/4.8 mL/1.2 mL), degassed with argon, and treated with Pd(OH)<sub>2</sub> (2.74 g). The mixture was placed under 1 atm of  
10 H<sub>2</sub>. After stirring for 18 h, the mixture was diluted with EtOAc and filtered through Celite. The filtrate was concentrated to give ester 13-4 as an off-white solid.  
TLC R<sub>f</sub> = 0.10 (5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>).  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.34 (s, 1H), 7.26 (dd, J=1.2, 8.1 Hz, 1H),  
15 7.12 (d, J=8.1 Hz, 1H), 4.65 (t, J=7.2 Hz, 1H), 4.13 (q, J=6.9 Hz, 2H), 2.98 (m, 2H), 1.20 (t, J=7.2 Hz, 3H) ppm.

3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl) propionic acid (13-5)

20 The title compound 13-5 was prepared from 2-9A and 13-4 using the procedure described in Scheme 2.  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.57 (d, J=7.3 Hz, 1H), 7.28 (s, 1H), 7.19 (d, J=8.2 Hz, 1H), 6.63 (d, J= 7.3 Hz, 1H), 5.47 (m, 1H), 3.30 (m, 9H), 2.82 (m, 2H), 2.66 (m, 2H), 1.96 (m, 6H) ppm.

25

## SCHEME 14

3-(4-Hydroxy-3-fluorophenyl)-acrylic acid ethyl ester (14-2)

A solution of 2-fluoro-4-bromophenol **14-1** (50 g, 261.8 mmol),  
 5 ethyl acrylate (34 mL),  $\text{Pd}(\text{OAc})_2$  (2.5 g), tri-*o*-tolylphosphine (5 g) and

triethylamine (83 mL) was heated to 100°C in a sealed tube for 12 h. The reaction was cooled to room temperature and diluted with dichloromethane (100 mL). The organic solution was washed with 10% aq. citric acid (40 mL), satd aq NaHCO<sub>3</sub>, and brine (40 mL). The organic solution was dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (50:50 hexanes/EtOAc to 100% EtOAc) to give acrylic acid 14-2 as a white solid. TLC R<sub>f</sub> = 0.45 (50% ethyl acetate/hexanes).

10 3-[Benzyl-(1(R)-phenylethyl)-aminol-3-(4-ethoxy-3-fluorophenyl)-propionic acid ethyl ester (14-4)

To a stirred solution of 14-2 (49.25 gm, 234.5 mmol) in DMF (600 mL) was added Cs<sub>2</sub>CO<sub>3</sub> (84.1 gm, 257.9 mmol) and ethyl iodide (18.8 mL, 234.5 mmol). After stirring for 12 h at room temperature, the reaction mixture was diluted with EtOAc (1L) and washed with water (6 x 300 mL), 10% aq. citric acid (200 mL), satd. aq. NaHCO<sub>3</sub> (200 mL), and brine (300 mL). The organic solution was dried over MgSO<sub>4</sub>, filtered, and concentrated to give 52.9 g (95%) of the product 14-3 as an orange oil which crystallized upon standing. A cooled (0°C) solution of (R)-(+)-N-benzyl- $\alpha$ -methylbenzylamine (71 mL, 339.4 mmol) in THF (650 mL) was treated with n-butyllithium (212 mL of a 1.6 M soln in hexanes; 339.4 mmol). After stirring for 10 min, the purple solution was cooled to -78°C and treated with a solution of ester 14-3 (53.8 g, 226.3 mmol) in THF (100 mL). After stirring for 20 min, the solution was quenched with satd aq NH<sub>4</sub>Cl soln (50 mL), and the cold bath removed. The reaction mixture was diluted with Et<sub>2</sub>O (1000 mL), and washed with 10% aq citric acid (300 mL), satd aq NaHCO<sub>3</sub> (300 mL), 5% aq acetic acid (300 mL), 10% aq K<sub>2</sub>CO<sub>3</sub> (300 mL), and brine (200 mL). The solution was dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (85:15 hexanes/EtOAc) to give the adduct 14-4. TLC R<sub>f</sub> = 0.39 (25% ethyl acetate/hexanes).

3-Amino-3-(4-Ethoxy-3-fluorophenyl)-propionic acid ethyl ester (14-5)

A solution of the dibenzylamine 14-4 (30.0 gm, 66.8 mmol) in EtOH/HOAc (340/30 mL) was purged with argon and treated with

Pd(OH)<sub>2</sub> (6 g) and placed under 1 atm of H<sub>2</sub> for 12 h. Additional portions (2.5 g) of Pd(OH)<sub>2</sub> were added after 24 h and 48 h. The reaction mixture was purged with argon, filtered through Celite, and the filtrate collected. The filtrate was concentrated to yield the desired amine 14-5.


5    <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.19 (m, 3H), 4.62 (m, 1H), 4.07 (m, 4H), 2.99 (m, 2H), 1.39 (m, 3H) 1.18 (m, 3H) ppm.

3-(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl) propionic acid (14-6)

10                    The title compound 14-6 was prepared from 2-9A and 14-5 using the procedure described in Scheme 2.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.45 (d, J = 7.3 Hz, 1H), 7.04 (m, 3H), 6.53 (d, J=7.3 Hz, 1H), 5.43 (m, 1H), 4.06 (q, J= 7.0 Hz, 2H), 3.48 (m, 6H), 3.15 (m, 1H), 2.78 (m, 6H), 2.55 (m, 2H), 1.96 (m, 3H), 1.38 (t, J= 7.0 Hz, 3H)  
15    ppm.

## SCHEME 15



5-Ethoxy-nicotinic acid ethyl ester (15-2)

A mixture of 3-hydroxy-nicotinic acid methyl ester 15-1 (15 g, 90.8 mmol), ethyl iodide (14.5 ml, 181.6 mmol), cesium carbonate (29.5 g, 90.8 mmol) and DMF (150 mL) was stirred at ambient temperature for 3 hours. The reaction mixture was diluted with Et<sub>2</sub>O and then washed with 10% K<sub>2</sub>CO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), and concentrated to give the ester 15-2 as a red oil.

TLC R<sub>f</sub> = 0.52 (silica, 75% EtOAc/hexanes)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.82 (s, 1H), 8.46 (s, 1H), 7.75 (s, 1H), 4.40 (q, 2H, J=7Hz), 4.12 (q, 2H, J=7Hz), 1.43 (m, 6H).

5-Ethoxy-N-methoxy-N-methyl-nicotinamide (15-3)

To a solution of 15-2 (15 g, 72 mmol) in EtOH (100 mL) was added 1N NaOH (80 ml, 80 mmol). After stirring for 1 h, the solvents were evaporated and the residue was dissolved in 1N HCl (80 ml, 80 mmol) and then concentrated, azeotroped with CH<sub>3</sub>CN to give the crude acid. The crude acid was suspended in DMF (200 mL) and then treated with HCl•HN(Me)OMe (13.9 g, 144 mmol), EDC (15.1g, 79.2 mmol), HOBT (9.6g, 72 mmol) and NMM (60 mL, 576 mmol). The mixture was stirred for 18 hours and then concentrated. The residue was dissolved in ethyl acetate, washed with 10% K<sub>2</sub>CO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>), and concentrated to give amide 15-3 as a brown oil.

TLC R<sub>f</sub> = 0.30 (silica, 70:25:5 chloroform/ ethyl acetate/ MeOH)

5-Ethoxy-pyridine-3-carbaldehyde (15-4)

To a stirred solution of 15-3 (14.0 g, 66.5 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (200 mL) at -78°C under argon was added DIBAL (1.0M hexanes, 90ml) dropwise over 30 minutes. After 30 minutes, the solution was warmed to 0°C for 1 hour. The reaction was quenched with 100 ml 1.0M Rochelle's salt, stirred for 1.0 hour and then extracted with Et<sub>2</sub>O. The organic layer was dried (MgSO<sub>4</sub>), and then concentrated to give the aldehyde 15-4 as a brown oil.

TLC R<sub>f</sub> = 0.32 (silica, 70:25:5 chloroform/ethyl acetate/MeOH)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.10 (s, 1H), 8.65 (s, 1H), 8.55 (s, 1H), 7.59 (s, 1H), 4.14 (q, 2H, J=7Hz), 1.43 (t, 3H, J=7Hz).

3-(5-Ethoxy-pyridin-3-yl)-acrylic acid *tert*-butyl ester (15-6)

A mixture of 15-4 (8.0 g, 51.6 mmol), 15-5 (20 g, 54.2 mmol), and benzene (150 mL) was heated to reflux for 30 minutes. The mixture was diluted with Et<sub>2</sub>O and then washed with 10% K<sub>2</sub>CO<sub>3</sub>, brine and  
5 dried (MgSO<sub>4</sub>). Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 30% EtOAc/hexanes) to give 15-6 as a yellow solid.

TLC R<sub>f</sub> = 0.41 (silica, 70:25:5 chloroform/ ethyl acetate/ MeOH)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.31 (m, 2H), 7.55 (d, 1H, J=16Hz), 7.27 (s,  
10 1H), 6.40 (d, 1H, J=16Hz), 4.10 (q, 2H, J=7Hz), 1.54 (s, 9H), 1.44 (m, 3H).

3(S)-Amino-3-(5-ethoxy-pyridin-3-yl)-propionic acid *tert*-butyl ester (15-8)

To a stirred solution of 15-7 (500mg, 2.38mmol) and THF at 0°C was added nBuLi (2.5 M THF, 0.95 ml) dropwise. After 20 minutes,  
15 the solution was cooled to -78°C and 15-6 (500mg, 1.98 mmol), dissolved in 3 ml THF, was added. After 15 minutes, the reaction was quenched with sat. NH<sub>4</sub>Cl followed by the removal of the cooling bath. The solution was extracted with ethyl acetate. The organic portion was washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The residue was dissolved in  
20 acetic acid (14 ml), and the solution was purged with argon for 30 minutes. 10% Pd/C (1.0 g) was added and the mixture was heated to 80°C. 1,4-Cyclohexadiene (6 ml) was added dropwise maintaining an internal temperature between 80°C and 90°C. After 5.0 hours, the mixture was filtered through a celite pad, concentrated and then  
25 azeotroped with toluene. The residue was chromatographed (silica gel, 5% [10:10:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O]/ 70:25:5 chloroform/ ethyl acetate/ MeOH) to give 15-8 as a yellow solid.


<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.18 (m, 2H), 7.25 (s, 1H), 4.41 (m, 1H), 4.08  
(q, 2H, J=7Hz), 2.59 (m, 2H), 1.87 (s, 2H), 1.40 (m, 12H).

3(S)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (15-9)

The title compound 15-9 was prepared from 2-9A and 15-8 using the procedure described in Scheme 2.

35 TLC R<sub>f</sub> = 0.27 (silica, 10:10:1:1 ethyl acetate/EtOH/water/NH<sub>4</sub>OH).

$^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.13 (m, 2H), 7.48 (d, 1H,  $J=7\text{Hz}$ ), 7.35 (s, 1H), 6.55 (d,  $J=8\text{ Hz}$ , 1H), 5.53 (m, 1H), 4.13 (q, 2H, 7Hz), 3.31-3.70 (m, 7H), 3.06 (m, 2H), 2.55 -2.85 (m, 6H), 1.88-2.15 (m, 5H), 1.42 (t, 3H,  $J=7\text{ Hz}$ ).

**SCHEME 16**



3(S)-Amino-3-(5-methoxy-pyridin-3-yl)-propionic acid tert-butyl ester  
(16-2)

3-Bromo-5-methoxy-pyridine 16-1 (prepared as described in *J. Org. Chem.* 1990, 55, 69) was converted into 16-2 utilizing the  
5 procedure previously described for the conversion of 1-4 to 1-6.  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.20 (d, 1H, J=3Hz), 8.18 (d, 1H,  
J=2Hz), 7.50 (s, 1H), 4.51 (m, 1H), 3.90 (s, 3H), 2.87 (m, 2H, ), 1.37 (m,  
9H).

10 3(S)-(5-Methoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl  
ester (16-3)

The title compound 16-3 was prepared from 2-9A and 16-2  
using the procedure described in Scheme 2.  
15 TLC R<sub>f</sub> = 0.27 (silica, 70:20:10 chloroform/ethyl acetate/MeOH)  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.23 (d, 1H, J=3Hz), 8.15 (s, 1H), 7.22 (s, 1H),  
7.02 (d, 1H, J=7Hz), 6.33 (d, 1H, 7Hz), 5.46 (t, 1H, J=8Hz), 4.78 (s, 1H),  
4.11 (m, 2H), 3.84 (s, 3H), 3.30 (m, 6H), 3.00 (m, 2H), 2.67 (t, 2H, J=6Hz),  
2.52 (m, 2H), 1.85 (m, 6H), 1.23 (m, 3H).

20 3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl  
ester (16-4)

To a stirred solution of 16-3 (200 mg, 0.4278mmol) and  
25 ethanethiol (0.5ml) and CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was added AlCl<sub>3</sub> (570mg, 4.28  
mmol). After 1.0 hour, the reaction was quenched with sat. NaHCO<sub>3</sub>.  
Ethyl acetate was added, and the reaction mixture was then purged with  
argon for 1.0 hour. The organic layer was separated, washed with  
brine, dried (MgSO<sub>4</sub>) and then concentrated to give the phenol 16-4 as a  
30 yellow oil.  
TLC R<sub>f</sub> = 0.22 (silica, 70:20:10 chloroform/ethyl acetate/MeOH)


3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1.8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (16-5)

The title compound 16-5 was prepared from 16-4 by basic hydrolysis using the procedure described in Scheme 2.

5 TLC R<sub>f</sub> = 0.39 (silica, 10:1:1 EtOH/water/NH<sub>4</sub>OH).

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.01 (m, 2H), 7.46 (d, 1H, J=7Hz), 7.20 (s, 1H), 6.53 (d, J=8 Hz, 1H), 5.49 (m, 1H), 3.51-3.68 (m, 2H), 3.46 (t, 2H, 5Hz), 3.19 (m, 2H), 3.00 (m, 2H), 2.52-2.78 (m, 6H), 1.92 (m, 4H).

SCHEME 17



3(S)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1.8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (17-2)


15 The title compound 17-2 was prepared from 2-9A and 17-1 (for preparation see J.A. Zablocki, et.al., *J. Med. Chem.* 1995, 38, 2378-2394) using the procedure described in Scheme 2.

TLC R<sub>f</sub> = 0.32 (silica, 15:10:1:1 ethyl acetate/EtOH/water/NH<sub>4</sub>OH).

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.45 (d, J=7Hz, 1H), 6.53 (d, J=8 Hz, 1H), 5.15 (m, 1H), 3.31-3.70 (m, 7H), 2.55-2.85 (m, 7H), 2.35 (m, 1H), 1.88 -2.15 (m, 4H).

20

SCHEME 18




2(S)-Benzenesulfonylamino-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (18-2)

The title compound 18-2 was prepared from 2-9A and 18-1  
 (for preparation, see Scheme A substituting phenylsulfonyl chloride for  
 5 4-iodophenylsulfonyl chloride) using the procedure described in Scheme  
 2.

TLC R<sub>f</sub> = 0.23 (silica, 15:10:1:1 ethyl acetate/EtOH/water/NH<sub>4</sub>OH).

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.81 (m, 2H), 7.36 (m, 3H), 7.10 (d, 1H,  
 J=8Hz), 6.37 (d, 1H, J=7Hz), 3.61 (m, 1H), 3.36 (m, 2H), 3.02 Æ 3.18 (m,  
 10 6H), 3.00 (m, 2H), 2.68 (t, 2H, J=6Hz), 2.50 (m, 2H), 1.79 Æ 1.90 (m, 4H).

SCHEME 19



3-Amino-pent-4-enoic acid ethyl ester (19-1)

15 A mixture of 5% Pd/BaSO<sub>4</sub> (0.025 g) and quinoline (0.30 mL)  
 was stirred under a balloon of hydrogen for 30 minutes. 3-Amino-pent-4-  
 ynoic acid ethyl ester 17-1 (1.77 g, 10.0 mmol) in EtOH (15 mL) was added  
 and the solution stirred for an additional 2.5 hours. The solution was  
 filtered through a pad of celite and concentrated in vacuo to provide 2.65  
 20 g of crude product 19-1.


<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.40-7.60 (br s, 2H), 6.11-5.96 (m, 1H), 5.58-5.53 (d, 1H), 5.44-5.41 (d, 1H), 4.31-4.16 (m, 3H), 3.12-2.86 (m, 2 H), 1.29-1.25 t, 3H).

5    3(S)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid (19-2)


The title compound 19-2 was prepared from 2-9A and 19-1 using the procedure described in Scheme 2.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 11.1 (s, 1H), 7.21-7.19 (d, 1H), 6.26-6.23 (d, 1H), 5.91-5.78 (m, 1H), 5.22-5.00 (m, 3H), 3.79-3.16 (m, 10H), 2.77-2.33 (m, 5H), 2.06-1.80 (m, 4H).  
10    MS (FAB) 359 (M+1)

## SCHEME 20



## SCHEME 20 (CONTINUED)



[3-(6-Bromo-pyridin-2-yl)-propyl]-(2-oxo-ethyl)-carbamic acid *tert*-butyl ester (20-2)

A solution of 2,6-dibromopyridine 20-1 (111 g, 468 mmol) and N-BOC-propargylamine (80.0 g, 515 mmol) in 500 ml of triethylamine at 0°C was treated with copper(I) iodide (2.23 g, 11.7 mmol). The mixture was purged with argon and then dichlorobis(triphenylphosphine)-palladium(II) (8.22 g, 11.7 mmol) was added. The solution was stirred at 0°C for one hour, then at room temperature for 16 hours. The solution was diluted with 250 mL ether and washed with H<sub>2</sub>O (4 X 100 mL). The organic extract was washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The solvents were removed in vacuo and the crude product was purified by silica gel chromatography (25% EtOAc/hexane) to afford 20-2.  
1H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.53-7.34 (m, 3 H), 4.82-4.80 (br s, 1 H), 4.18-4.17 (d, 2 H), 1.46 s, 9 H).

3-(2-([3-(6-Bromo-pyridin-2-yl)-propyl]-*tert*-butoxycarbonyl-amino)-ethylamino) (20-3)

To a solution of 20-2 (79.8 g, 257 mmol) in 350 mL of ethanol and triethylamine (26.8 mL, 193 mmol) was added platinum(IV) oxide (2.91 g, 12.8 mmol). After stirring under a hydrogen atmosphere for 4 hours, the solution was filtered through a pad of celite and concentrated in vacuo. The crude product was dissolved in EtOAc (200 mL) and washed with H<sub>2</sub>O (4 X 250 mL) and brine (250 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated in vacuo. The crude material was purified by silica gel chromatography (10% EtOAc/CHCl<sub>3</sub>) to afford 20-3.  
1H NMR (CDCl<sub>3</sub>, 400 MHz): δ 7.48-7.42 (t, 1 H), 7.32-7.29 (d, 1H), 7.13-7.10 (d, 1H), 4.71-4.70 (br s, 1 H), 3.18-3.09 (m, 2 H), 2.82-2.77 (t, 2 H), 1.96-1.85(m, 2 H), 1.44 (s, 9 H).

3-(6-Bromo-pyridin-2-yl)-propylamine•hydrochloride (20-4)

A solution of 20-3 (3.33 g, 10.5 mmol) in EtOAc (150 mL) was saturated with HCl gas and stirred at room temperature for 2 hours. The solvent was removed in vacuo to provide 20-4. The crude product was used in the next step without further purification.

[3-(6-Bromo-pyridin-2-yl)-propylamino]-acetic acid ethyl ester (20-5)

A solution of 3-(6-bromo-pyridin-2-yl)-propylamine 20-4 (25.6 g, 89.1 mmol), diisopropylethylamine (46.5 mL, 267 mmol), acetic acid (28 mL, 490 mmol), and ethyl glyoxylate (10.9 g, 107 mmol) in 200 mL of methanol was stirred at room temperature for one hour. A 1M solution of NaCNBH<sub>3</sub> in THF (98.0 mL, 98.0 mmol) was added slowly with a syringe pump over 4 hours. The resulting solution was stirred for 12 hours, after which the solvent was removed in vacuo and the residue taken up in chloroform and filtered. The solution was then washed with 10% Na<sub>2</sub>CO<sub>3</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvent removed in vacuo to give the crude amine. The crude product was purified by silica gel chromatography (7% MeOH/CHCl<sub>3</sub>) to give 20-5 in a 3:2 mixture of ethyl and methyl esters.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.47-7.42 (t, 1H), 7.31-7.27, (t, 1H), 7.13-7.10 (d, 1H), 4.20-4.14, (m, 2H), 3.39 (s, 2H), 2.85-2.75 (m, 2H), 2.68-2.63 (t, 2H), 1.96-1.88 (m, 2H), 1.29-1.24 (m, 3H).

[[3-(6-Bromo-pyridin-2-yl)-propyl]-tert-butoxycarbonyl-amino]-acetic acid ethyl ester (20-6)

To a solution of [3-(6-bromo-pyridin-2-yl)-propylamino]-acetic acid ethyl ester 20-5 (17.6 g, 58.6 mmol) in THF (200 mL) was added di-*tert*butyldicarbonate (15.3 g, 70.3 mmol). After stirring at room temperature for 16 hours, the solvents were removed in vacuo. The product was purified by silica gel chromatography (5% MeOH/CHCl<sub>3</sub>) to give 20-6.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.47-7.42 (m, 1H), 7.32-7.28 (t, 1H), 7.16-7.10 (t, 1H), 4.22-4.15, (q, 2H), 3.95-3.85 (d, 2H), 3.38-3.29 (m, 2H), 2.80-2.75 (t, 2H), 2.03-1.91 (m, 2H), 1.46-1.44 (m, 9H), 1.31-1.23 (m, 3H).

[3-(6-Bromo-pyridin-2-yl)-propyl]-[(methoxy-methyl-carbamoyl)-methyl]-carbamic acid *tert*-butyl ester (20-7)

To a solution of 20-6 (23.4 g, 58.4 mmol) in ethanol (200 mL) was added NaOH (100 mL 1M solution in water, 100 mmol). After stirring for 1 h at 50°C, HCl (10.3 mL 12 M, 4.75 mmol) was added in 50 mL EtOH, and the mixture evaporated to give an oily residue. The



residue was evaporated from ethanol three times, and then from acetonitrile three times, producing a yellow crusty solid which was dried under a vacuum of <2 mm Hg for 2 h. This residue was then slurried in acetonitrile (180 mL) and chloroform (180 mL), and NMM (41.7 mL, 379 mmol), N,O-dimethylhydroxylamine hydrochloride (11.9 g, 122 mmol), HOBT (10.2 g, 75.9 mmol), and EDC (14.5 g, 75.9 mmol) were added. After stirring for 15 h, the mixture was evaporated to dryness, the residue slurried in EtOAc, washed with sat. NaHCO<sub>3</sub>, brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporative removal of the solvent followed by evaporation from toluene to remove the residual NMM gave 20-7 as a yellow oil. TLC R<sub>f</sub> = 0.49 (silica, 70:25:5 chloroform/ethyl acetate/methanol) <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.48-7.41 (m, 1H), 7.32-7.28 (t, 1H), 7.17-7.11 (t, 1H), 4.14 (s, 2H), 3.73-3.70 (d, 3H), 3.39-3.30 (m, 2H), 3.18 (s, 3H), 2.80-2.75 (t, 2H), 2.02-1.91 (m, 2H), 1.45 (m, 9H).

[3-(6-Bromo-pyridin-2-yl)-propyl]-(2-oxo-ethyl)-carbamic acid tert-butyl ester (20-8)

To a stirred solution of 20-7 (14.9 g, 35.7 mmol) and THF (100 ml) at -78°C was added DIBAL (1.0M/hexanes, 53.6 ml, 53.6 mmol) dropwise over 20 minutes. After 1 h, the mixture was warmed to RT and quenched by the careful addition of 20 mL MeOH. 200 ml of 1.0 M Rochelle's salt was then added followed by the removal of the cooling bath. The mixture was stirred for 1.0 hour and then diluted with Et<sub>2</sub>O. After another 30 minutes of stirring, the organic portion was separated and dried over MgSO<sub>4</sub>. Evaporative removal of the solvent gave the crude aldehyde 20-8 as a colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 9.59-9.56 (d, 1H), 7.48-7.43 (t, 1H), 7.32-7.26 (m, 1H), 7.14-7.07 (m, 1H), 3.53-3.26 (m, 4H), 2.80-2.72 (m, 2H), 2.00-1.93 (m, 2H), 1.44 (s, 9H).

3-(5-Ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester (20-9)

A mixture of the crude aldehyde 20-8 (0.671 g, 1.88 mmol), amine 15-8 (0.651 g, 2.44 mmol), acetic acid (0.107 mL, 1.88 mmol), NaOAc (1.54 g, 18.8 mmol), and powdered molecular sieves (1.20g) in 2-propanol (15 mL) was stirred for 20 minutes. The mixture was cooled to

0°C and then NaBH<sub>3</sub>CN (0.354 g, 5.64 mmol) was added. After stirring 6 hours, the pH of the mixture was adjusted to ~2 with 1N HCl. The solution was stirred for an additional 10 minutes, ethyl acetate (20 mL) was added, and the pH was adjusted to ~11 with 10% potassium carbonate. The organics were extracted with ethyl acetate, dried over Na<sub>2</sub>SO<sub>4</sub>, and removed in vacuo. The residue was chromatographed (silica gel, [70:25:5 CHCl<sub>3</sub>/EtOAc/MeOH]) to give 20-9 as a yellow oil in 95% yield.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.19-8.18 (d, 1H), 8.14-8.13 (d, 1H), 7.48-7.42 (t, 1H), 7.32-7.26 (m, 1H), 7.23-7.20 (m, 1H), 7.11-7.08 (m, 1H), 4.16-4.02 (m, 3H), 3.30-3.18 (m, 4H), 2.74-2.45 (m, 7H), 1.92-1.86 (t, 2H), 1.45-1.38 (m, 21H).

3-(2-[3-(6-Bromo-pyridin-2-yl)-propylamino]-ethylamino)-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester (20-10)

To a solution of 3-(5-ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester 20-9 (0.085g, 0.141 mmol) in dichloromethane (5 mL) was added *p*-toluenesulfonic acid (0.161 g, 0.847 mmol). The mixture was stirred for 2 hours at room temperature and was then neutralized with 1N NaOH. The organic layer was extracted (3 X 25 mL) with CHCl<sub>3</sub>, dried, and concentrated in vacuo. The crude product 20-10 was not purified (0.069 g, 96% yield).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.23-8.21 (m, 2H), 7.45 (s, 1H), 7.40-7.35 (m, 1H), 7.28-7.25 (t, 1H), 7.04-7.01 (d, 1H), 6.28 (br s, 2H) 4.39-4.33 (t, 1H), 4.00-3.92 (q, 2H), 3.40-3.35 (m, 2H), 3.28-3.22 (m, 1H), 3.15-2.90 (m, 4H), 2.79-2.71 (m, 3H), 2.14-2.01 (m, 2H), 1.34-1.26 (m, 12H).

3-(3-[3-(6-Bromo-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl)-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester (20-11)

To a stirred solution of 20-10 (0.80 g, 1.57 mmol) and diisopropylethylamine (0.823 mL, 4.72 mmol) in dichloromethane (10 mL) at 0°C was added *p*-nitrophenyl chloroformate (0.333 g, 1.65 mmol). The solution stirred for 30 minutes and dioxane (10 mL) was added, then refluxed for 4 hours. EtOAc (100 mL) was added and the organics were

washed with 10% K<sub>2</sub>CO<sub>3</sub>, dried, and concentrated in vacuo. The residue was chromatographed (silica gel, [70:20:10 CHCl<sub>3</sub>/ EtOAc/ MeOH]) to give 20-11.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.22-8.18 (dd, 1H), 8.14-8.13 (t, 1H), 7.46-7.37 (m, 1H), 7.31-7.24 (m, 1H), 7.20-7.16 (m, 1H), 7.11-7.09 (d, 1H), 4.09-4.04 (q, 2H), 3.34-3.16 (m, 5H), 2.99-2.87 (m, 2H), 2.77-2.69 (m, 2H), 2.63-2.46 (m, 2H), 1.97-1.88 (m, 2H), 1.44-1.37 (m, 12H).

MS M+1 = 533.3

10 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-{3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl}-2-oxo-imidazolidin-1-yl)-propionic acid *tert*-butyl ester (20-12)

To a stirred solution of 20-11 (0.075 g, 0.142 mmol) in toluene (3 mL) was added Pd(DBA)<sub>2</sub> (0.0041 g, 0.0071 mmol), DPPF (0.0039 g, 0.0071 mmol), and NaOt-Bu (0.0163 g, 0.170 mmol) followed by *p*-methoxybenzylamine (0.0204 mL, 0.156 mmol). The resulting solution was heated at 110°C for 2 hours. The solution was cooled and the solvent was removed in vacuo. The product was purified by silica gel chromatography (10% EtOH/EtOAc) to give 20-12.

15 <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.21-8.20 (d, 1H), 8.14-8.13 (d, 1H), 7.33-7.25 (m, 3H), 7.19-7.17 (t, 1H), 6.87-6.84 (d, 2H), 6.45-6.43 (d, 1H), 6.21-6.18 (d, 1H), 5.48-5.42 (t, 1H), 5.26 (br s, 1H), 4.38-4.37 (d, 2H), 4.09-4.01 (q, 2H), 3.79 (s, 3H), 3.31-3.18 (m, 5H), 3.07-2.87 (m, 3H), 2.63-2.58 (t, 2H), 1.95-1.84 (m, 2H), 1.49-1.39 (m, 12H).

25 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-{3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl}-2-oxo-imidazolidin-1-yl)-propionic acid (20-13)

To a stirred solution of 20-12 (0.028 g, 0.047 mmol) in dichloromethane (10 mL) was added TFA (1 mL). After 1 hour, the solvent was removed in vacuo and azeotroped twice with toluene (15 mL). The residue was chromatographed (silica gel, 25:10:1:1 followed by 15:10:1:1 ethyl acetate /EtOH /water/NH<sub>4</sub>OH) to give 20-13 as a white solid.

30 <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz): δ 8.21-8.20 (d, 2H), 7.50-7.46 (t, 1H), 7.31-7.26 (m, 2H), 7.19 (s, 1H), 6.88-6.85 (d, 2H), 6.42-6.40 (d, 1H), 6.34-6.32 (d, 1H),

5.69-5.63 (m, 1H), 5.30 (s, 1H), 4.43-4.41 (d, 2H), 4.09-4.01 (q, 2H), 3.77 (s, 3H), 3.75-3.44 (m, 3H), 3.24-2.86 (m, 4H), 2.79-2.67 (m, 3H), 2.03-1.90 (m, 2H), 1.44-1.41 (t, 3H).

MS (FAB) 534 (M+1)

5


3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid (20-14)



To a stirred solution of 20-13 (0.031 g, 0.052 mmol) in dichloromethane (10 mL) was added TFA (1 mL). The solution was stirred for 16 hours at 85°C. after which the solvent was removed in vacuo and azeotroped twice with toluene (15 mL). The residue was chromatographed (silica gel, 15:10:1:1 followed by 10:10:1:1 ethyl acetate /EtOH /water /NH<sub>4</sub>OH) to give 20-14 as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 400 MHz): δ 8.13-8.10 (m, 2H), 7.59-7.54 (t, 1H), 7.38-7.35 (m, 1H), 6.60-6.57 (d, 2H), 5.53-5.47 (q, 1H), 4.15-4.09 (q, 2H), 3.64-3.57 (m, 1H), 3.47-3.41 (m, 1H), 3.28-3.21 (m, 2H), 3.07-2.90 (m, 3H), 2.76-2.61 (m, 3H), 2.02-1.83 (m, 2H), 1.42-1.38 (t, 3H).

MS (FAB) 414 (M+1)

## Scheme 21



SCHEME 21 (CONTINUED)3-Bromo-6-chloro-5-nitropyridine (21-2)

- A suspension of  $\text{CuCl}_2$  (3.33 g, 24.8 mmol) in anhydrous  $\text{CH}_3\text{CN}$  (200 mL) at  $65^\circ$  was treated with tert-butyl nitrite (3.13 mL, 26.3 mmol), followed by the dropwise addition of a solution of 21-1 in 60 mL of  $\text{CH}_3\text{CN}$ . The resulting mixture was stirred under an argon atmosphere at  $65^\circ$  for 2 h and concentrated at reduced pressure. The residue was partitioned between EtOAc (150 mL) and 3% HCl (60 mL), and the organic layer washed successively with 3% HCl, water, and brine (60 mL), then dried, filtered and concentrated to afford a brown solid which was chromatographed on silica (25% EtOAc/Hexane) to afford 21-2 as a yellow crystalline solid.
- TLC  $R_f$  = 0.60 (25% EtOAc/ Hexane)
- $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.70 (d,  $J=2.4$  Hz, 1H), 8.37 (d,  $J=2.4$  Hz, 1H).

(3-Nitro-5-bromo-pyridin-2-yloxy)-acetic acid methyl ester (21-3)

Methyl glycolate (450 mg, 5.05 mmol) was added to a suspension of 60% NaH (131 mg, 55 mmol) in THF (20 mL) at 0°. The resulting solution was stirred under argon for 0.5 h, then treated with a solution of 21-2. After stirring at 0° for 0.5 h, the reaction was diluted with ethyl acetate, and washed successively with sat. NaHCO<sub>3</sub>, water and brine (80 mL each), then dried, filtered and concentrated to afford 21-3 as a yellow solid.

TLC R<sub>f</sub> = 0.70 (25% EtOAc/ Hexane)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.46 (d, J = 2.4 Hz, 1H), 8.37 (d, J = 2.4 Hz, 1H) 5.15 (s, 2H), 3.78 (s, 3H).

2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-7-bromo-naphthalene (21-4)

A mixture of 21-3 (1.5 g, 5.12 mmol) and powdered tin (1.37 g, 11.5 mmol) was treated with conc. HCl (10 mL). The mixture was heated to 80° for 2 h, then cooled and concentrated. The residue was partitioned between CHCl<sub>3</sub> and sat. NaHCO<sub>3</sub>, washed with brine, then dried, filtered and concentrated to afford a yellow solid.

Chromatography on silica gel (50% hexane/EtOAc) gave 21-4 as a yellow solid.

TLC R<sub>f</sub> = 0.65 (50% EtOAc/ Hexane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 10.81 (br,s, 1H), 7.88 (d, J=2.4 Hz, 1H), 7.25 (d, J=2.4 Hz, 1H), 4.81 (s, 2H).

3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-acrylic acid tert-butyl ester (21-5).

A mixture of 21-4 (1.12 g, 4.89 mmol), (o-tol)<sub>3</sub>P (298 mg, 1.0 mmol), Pd(OAc)<sub>2</sub> (110 mg, 0.49 mmol), and triethylamine (0.86 mL, 5.87 mmol) in DMF (20 mL) was placed in a 100-mL flask. The mixture was degassed with argon, then tert-butyl acrylate (752 mg, 5.87 mmol) was added and the tube sealed and heated to 100° for 12 h. The reaction mixture was diluted with ethyl acetate, filtered and washed with NaHCO<sub>3</sub>, water, and brine, dried, filtered and concentrated.

Chromatography on silica gel (25% hex/EtOAc) gave 21-5 as a yellow solid.

TLC R<sub>f</sub> = 0.60 (25% EtOAc/ Hexane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 10.91 (br,s, 1H), 8.15 (d, J=2.4 Hz, 1H), 7.54 (d, J=16 Hz, 1 H), 7.42 (d, J= 2.4 Hz, 1H), 6.35 (d, J=16 Hz, 1 H), 4.84 (s, 2H), 1.48 (s, 9H).

5    3(S)-[Benzyl-(1(R)-phenylethyl)-amino]-3-(2-oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert*-butyl ester (21-6)

A solution of N-benzyl-α-(R)-methylbenzylamine (0.82 g, 3.87 mmol) in THF (25 mL) at 0°C was treated with n-BuLi (1.6 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0°C for 30 min and then cooled to -78°C. A solution of acrylate 21-5 (0.485 g, 1.76 mmol) in THF (5 mL) was added. After stirring for 15 min at -78°C, satd aq NH<sub>4</sub>Cl soln (5 mL) was added and the cold bath removed. The mixture was warmed to room temperature, and extracted with Et<sub>2</sub>O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated. The residue was purified by flash chromatography (40% ethyl acetate/hexanes) to give the β-aminoester 21-6 as a yellow oil.

TLC R<sub>f</sub> = 0.3 (40% ethyl acetate/hexanes)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1H NMR 8.70 (br, s, 1H), 7.91 (d, J=1.8 Hz, 1H), 7.4-7.2 (10H), 7.12 (d, J=1.8 Hz, 1H), 4.80 (s, 2 H), 4.42 (m, 1H), 3.91 (q, J=6.7 Hz, 1 H), 3.69 (d, J=7.2 Hz, 1H, ), 3.62 (d, J =7.2 Hz, 1H, ), 2.46 (m, 2H), 1.34 (d, J=7.0 Hz, 3H), 1.29 (s, 9H).

25    3(S)-Amino-3-(2-oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert*-butyl ester (21-7)

A mixture of the dibenzylamine 21-6 (0.22 g, 0.44 mmol) in EtOH/H<sub>2</sub>O/AcOH (26 mL/3 mL/1.0 mL) was degassed with argon and treated with Pd(OH)<sub>2</sub> (100 mg). The mixture was placed under 1 atm of H<sub>2</sub>. After stirring for 18 h, the mixture was diluted with EtOAc and filtered through celite. The filtrate was concentrated and the residue purified by flash chromatography (20% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 80% EtOAc) to give the *tert*-butyl ester 21-7 as a white solid.

TLC R<sub>f</sub> = 0.5 (20% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 80% EtOAc)



<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.89 (d, J = 1.7 Hz, 1H), 7.31 (d, J = 1.7 Hz, 1H), 4.81 (s, 2H), 4.38 (m, 1H), 2.6, (m, 2H), 1.41 (s, 9H).

5 3(R)-[Benzyl-(1-phenylethyl)-aminol-3(S)-(2-thioxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert.* butyl ester (21-8)

A solution of 21-6 (0.22 g, 0.44 mmol) in anhydrous THF was treated with Lawesson's reagent (0.098 g, 0.243 mmol) and stirred at room temperature for 1.5 h. Silica gel (500 mg) was added to the reaction mixture and the solvent was removed at reduced pressure and the product was eluted from silica using 25% EtOAc/hexane to afford 21-8 as a yellow solid.

TLC R<sub>f</sub> = 0.7 (40% EtOAc/hexane)

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 9.82 (br, s, 1H), 7.95 (d, J = 1.8 Hz, 1H), 7.4-7.2 (11H), 5.08 (s, 2H), 4.42 (m, 1H), 3.91 (q, J = 6.7 Hz, 1H), 3.69 (d, J = 7.2 Hz, 1H), 3.62 (d, J = 7.2 Hz, 1H), 2.46 (m, 2H), 1.34 (d, J = 7.0 Hz, 3H), 1.29 (s, 9H).

3(S)-Amino-3-(2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert.* butyl ester (21-9)

20 A solution of 21-8 (1.0 g, 1.9 mmol) in anhydrous Et<sub>2</sub>O (10 mL) at 0° was treated dropwise with LiAlH<sub>4</sub> (2.09 ml of a 1.0 M solution in Et<sub>2</sub>O). The resulting solution was stirred at 0°C for 30 min and then quenched by the sequential addition of H<sub>2</sub>O (0.3 mL), 15 % NaOH (0.08 mL). Celite (1 g) was added and the mixture filtered through a Celite pad. The filtrate was evaporated and the residue was purified by flash chromatography (65% ethyl acetate/hexanes) to give the dibenzylamine intermediate as a yellow oil.

TLC R<sub>f</sub> = 0.4 (65% ethyl acetate/hexanes)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1H NMR 7.61 (d, J = 1.8 Hz, 1H), 7.4-7.2 (10H), 6.87 (d, J = 1.8 Hz, 1H), 4.41 (m, 2H), 4.36 (m, 1H), 3.91 (q, J = 6.7 Hz, 1H), 3.8 (brs, 1H), 3.69 (m, 2H), 3.42 (m, 2H), 2.46 (m, 2H), 1.34 (d, J = 7.0 Hz, 3H), 1.29 (s, 9H).

This material was deprotected with Pd(OH)<sub>2</sub> in ethanol to afford 21-9 as a white solid. TLC R<sub>f</sub> = 0.5 (20% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 80% EtOAc)

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.59 (d, J=1.7 Hz, 1H), 6.92 (d, J=1.7 Hz, 1H), 4.41 (m, 2H), 4.30 (m, 1H), , 3.41 (m, 2H), 2.6, (m, 2H), 1.41 (s, 9H).

5 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5.6.7.8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (21-10)

The title compound 21-10 was prepared from 2-9A and 21-7 using the procedure described in Scheme 2.

High resolution MS Calc'd.=418.2198, Obs'd = 481.2193.


10

3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5.6.7.8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (21-11)

15 The title compound 21-11 was prepared from 2-9A and 21-8 using the procedure described in Scheme 2.

High resolution MS Calc'd.=467.2417, Obs'd = 467.2401.

Scheme 22



**3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-7-bromo-naphthalene (22-2)**

A solution of **22-1** (4.8 g, 32 mmol) in MeOH (160 mL) at  $-15^\circ$  was treated dropwise with bromine (25.7 g, 161 mmol). After stirring at  $-15^\circ$  for 0.5 h, the mixture was warmed to ambient temperature and stirred overnight. The resulting white precipitate was filtered and washed with cold MeOH to afford **22-2** as a white solid.

TLC Rf = 0.65 (50% EtOAc/ Hexane)

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  11.2 (br s, 1H), 8.05 (d,  $J=2.4$  Hz, 1H), 7.66 (d,  $J=2.4$  Hz, 1H), 4.76 (s, 2H).

3(S)-Amino-3-(3-oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-propionic acid tert-butyl ester (22-3)

- 5 Bromide 22-2 was converted to amino ester 22-3 as illustrated in Scheme 21.  
TLC R<sub>f</sub> = 0.5 (12% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 88% EtOAc)  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.04 (d, J = 1.7 Hz, 1H), 7.34 (d, J = 1.7 Hz, 1H), 4.76 (s, 2H), 4.38 (m, 1H), 2.6, (m, 2H), 1.41 (s, 9H).

10 3(S)-Amino-3-(3-oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-propionic acid tert-butyl ester (22-4)

- Bromide 22-2 was converted to amino ester 22-4 as illustrated in Scheme 21.  
TLC R<sub>f</sub> = 0.5 (20% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 80% EtOAc)  
15 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.04 (d, J = 1.7 Hz, 1H), 7.34 (d, J = 1.7 Hz, 1H), 4.76 (s, 2H), 4.38 (m, 1H), 2.6, (m, 2H), 1.41 (s, 9H).


20 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (22-5)

The title compound 22-5 was prepared from 2-9A and 22-3 using the procedure described in Scheme 2.  
High resolution MS Calc'd.=481.2198, Obs'd = 481.2194.

25 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (22-6)

- The title compound 22-6 was prepared from 2-9A and 22-4 using the procedure described in Scheme 2.  
30 High resolution MS Calc'd.=467.2417, Obs'd = 467.2411.

Scheme 23

Furo-[2,3-b]pyridine-5-carboxaldehyde (23-2)

- 5 A solution of alcohol 23-1 (M. Bhupathy, et al., J. Heterocycl. Chem. 1995, 32, 1283-1287) was treated with excess  $\text{MnO}_2$  (10 eq) and the mixture stirred at room temperature for 16 h, then filtered through Celite and evaporated to afford 23-2 as a white solid. TLC  $R_f = 0.40$  (25% EtOAc/Hexane)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.22 (s, 1H), 9.05 (d, J= 1.8 Hz, 1H), 8.27 (d, J=1.7 Hz, 1H) 8.08 (d, J=1.8 Hz, 1H), 7.10 (d, J=1.7 Hz, 1H).

3-Amino-3-(furo[2.3-b]pyridin-5-yl)-propionic acid ethyl ester (23-3)

5 A solution containing aldehyde 23-2 (1.5 g, 10 mmol), ethyl hydrogen malonate (1.6 g, 20 mmol), and ammonium acetate (3.8 g, 50 mmol) in anhydrous ethanol (125 mL) was heated at reflux for 8 h. After cooling to room temperature, the solvent was evaporated and the residue partitioned between sat. sodium bicarbonate and EtOAc, the organic  
10 layer removed, dried, and concentrated. Chromatography of the residue afforded the amino ester 23-3 as a waxy solid.  
TLC R<sub>f</sub> = 0.5 (20% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 80% EtOAc)  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.34 (d, J=1.7 Hz, 1H), 8.04 (d, J=1.7 Hz, 1H), 7.72 (d, J= 1.7 Hz, 1H), 6.78 (d, J = 1.7 Hz, 1H), 4.62 (m, 1H), 4.13 (q,  
15 J=7.5 Hz, 2H), 3.20 (br, s, 2H), 2.76 (m, 2H), 1.23 (t, J= 7.5 Hz, 3H).


3-Furo[2.3-b]pyridin-6-yl-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1.8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid (23-4)

The title compound 23-4 was prepared from 2-9A and 23-3  
20 using the procedure described in Scheme 2.  
TLC R<sub>f</sub> = 0.30 (50% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O -50% EtOAc).  
FAB MS Obs'd 450.1 (M+H).

3-(2,3-Dihydrofuro[2.3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1.8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid (23-5)

25 A solution of 23-4 (360 mg, 0.80 mmol) in MeOH (10 mL) was treated with 10% Pd/C (100 mg) and stirred under a hydrogen atmosphere for 18h. The catalyst was removed by filtration through celite and the residue chromatographed (75% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O -  
30 25% EtOAc) to afford 23-5 as a colorless glass.  
TLC R<sub>f</sub> = 0.30 (50% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O -50% EtOAc)  
FAB MS Obs'd 452.2 (M+H).

Scheme 24

**Furo[3,2-b]pyridine-5-carboxaldehyde (24-2)**

- A solution of alcohol **24-1** (J.M. Hoffman, Jr., US Patent No. 4,808,595) was treated with excess  $\text{MnO}_2$  (10 eq) and the mixture stirred at room temperature for 16 h, then filtered through Celite and evaporated to afford **24-2** as a white solid.
- $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  10.18 (s, 1H), 8.92 (d,  $J=1.8$  Hz, 1H), 8.17 (d,  $J=1.7$  Hz, 1H) 7.89 (d,  $J=1.8$  Hz, 1H), 7.10 (d,  $J=1.7$  Hz, 1H).

3-Amino-3-(furo[3.2-b]pyridin-5-yl)-propionic acid ethyl ester (24-3)

- A solution containing aldehyde 24-2 (1.5 g, 10 mmol), ethyl hydrogen malonate (1.6 g, 20 mmol), and ammonium acetate (3.8 g, 50 mmol) in anhydrous ethanol (125 mL) was heated at reflux for 8 h. After cooling to room temperature, the solvent was evaporated and the residue partitioned between sat. sodium bicarbonate and EtOAc, the organic layer removed, dried, and concentrated. Chromatography of the residue afforded the amino ester 24-3 as a waxy solid.
- 10 TLC R<sub>f</sub> = 0.5 (20% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 80% EtOAc)  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.58 (d, J=1.7 Hz, 1H), 7.89 (d, J=1.7 Hz, 1H), 7.85 (d, J=1.7 Hz, 1H), 6.98 (d, J=1.7 Hz, 1H), 4.62 (t, J=7.2 Hz, 1H), 4.09 (q, J=7.5 Hz, 2H), 2.76 (m, 2H), 2.20 (br, s, 2H), 1.21 (t, J=7.5 Hz, 3H).

15 3-(Furo[3.2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1.8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid (24-4)


- The title compound 24-4 was prepared from 2-9A and 24-3 using the procedure described in Scheme 2.
- TLC R<sub>f</sub> = 0.56 (75% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 25% EtOAc)
- 20 High resolution MS Calc'd.=450.2117, Obs'd = 450.2136.

3-(2,3-Dihydrofuro[3.2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1.8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid (24-5)

- A solution of 24-4 (181 mg, 0.38 mmol) in acetic acid (5 mL) was treated with PtO<sub>2</sub> (100 mg) and stirred under a hydrogen atmosphere for 1h. The catalyst was removed by filtration through celite and the residue chromatographed (75% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 25% EtOAc) to afford 24-5 as a colorless glass.
- 25 TLC R<sub>f</sub> = 0.50 (75% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O - 25% EtOAc)
- 30 High resolution MS Calc'd.=452.2298, Obs'd = 452.2238



Scheme 25



N-(S)-(2-Amino-phenyl)-3-tert-butoxycarbonylamino-succinamic acid methyl ester (25-3a)

- 5                   A mixture of Boc-L-aspartic acid- $\beta$ -methyl ester 25-1 (5.0g, 20.2 mmol), o-phenylenediamine 25-2a (2.2 g, 20.2 mmol), EDC (3.9 g, 20.2 mmol), HOAT (0.28 g, 2.02 mmol), and NMM (6.7 mL, 60.7 mmol) in DMF (50 mL) was stirred for 18 h at ambient temperature. The solution was diluted with EtOAc (250 mL) and washed with sat. sodium
- 10 bicarbonate, water, and brine (50 mL each), then dried and evaporated to afford 25-3a as a yellow solid.
- TLC  $R_f$  = 0.50 (95%  $\text{CHCl}_3$ /5% isopropanol)
- $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.10 (br,s, 1H), 7.23 (d,  $J$ = 7.8 Hz, 1H), 7.08 (t,  $J$ =7.8 Hz, 1H) 6.78 (m, 1H), 5.8 (br d, 1H), 4.65 (m, 1 H), 3.76 (s, 3H),
- 15 3.15 (dd,  $J$ =4.6, 16 Hz, 1H), 2.90 (dd,  $J$ = 5.1, 16 Hz, 1H), 1.48 (s, 9H).

3(S)-Amino-3-(benzimidazol-2-yl)-propionic acid methyl ester (25-4a)

Ester 25-3a (1.0 g, 3 mmol) was dissolved in acetic acid (50 mL) and heated to 65° for 2 h. The solvent was removed to afford the Boc-protected intermediate as a white solid. The crude material (920 mg, 2.43 mmol) was dissolved in EtOAc, cooled to 0°, and treated with HCl gas to give 25-4a as a tan solid.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.80 (m, 2H), 7.35 (m, 2H), 5.98 (m, 1H), 3.80 (m, 2H), 3.76 (s, 3H).

10

3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (25-5a)

The title compound 25-5a was prepared from 2-9A and 25-4a using the procedure described in Scheme 2.

15 TLC R<sub>f</sub> = 0.30 (50% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O-50% EtOAc).

FAB MS Obs'd 449.2 (M+H).


3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (25-5b)

20 The title compound 25-4b was prepared as described above substituting 3,4-diaminopyridine for o-phenylenediamine.

TLC R<sub>f</sub> = 0.25 (50% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O -50% EtOAc).

FAB MS Obs'd 450.2 (M+H).

Scheme 26



N-(S)-(2-Hydroxy-phenyl)-3-tert-butoxycarbonylamino-succinamic acid methyl ester (26-2)

A mixture of Boc-L-aspartic acid-β-methyl ester (25-1) (5.0g, 20.2 mmol), 2-amino phenol (26-1) (2.2 g, 20.2 mmol), EDC (3.9 g, 20.2 mmol), HOAT (0.28 g, 2.02 mmol), and NMM (6.7 mL, 60.7 mmol) in DMF (50 mL) was stirred for 18 h at ambient temperature. The solution was diluted with EtOAc (250 mL) and washed with sat. sodium bicarbonate, water, and brine (50 mL each), then dried, and evaporated and chromatographed on silica (EtOAc) to afford 26-2 as a white solid. TLC R<sub>f</sub> = 0.55 (EtOAc)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.23 (d, J = 7.8 Hz, 1H), 6.89 (t, J = 7.8 Hz, 1H), 6.78 (m, 1H), 5.68 (br d, 1H), 4.65 (m, 1H), 3.76 (s, 3H), 3.15 (dd, J = 4.6, 16 Hz, 1H), 2.90 (dd, J = 5.1, 16 Hz, 1H), 1.48 (s, 9H).

3(S)-Amino-(3-benzoxazol-2-yl)-propionic acid methyl ester (26-3)


Ester 26-2 (2.0 g, 6.0 mmol) was dissolved in anhydrous THF (150 mL) along with Ph<sub>3</sub>P (1.58 g, 6.0 mmol). The resulting solution was cooled to 0°, and a solution of diethyl azodicarboxylate (1.53 g, 6.2 mmol) in THF (25 mL) was added dropwise. The cooling bath was removed and the solution stirred overnight at ambient temperature. The solution was concentrated and the residue chromatographed (75% EtOAc/Hexane) to afford the Boc-protected ester as a colorless glass. The crude material (1.8 g, 5.0 mmol) was dissolved in EtOAc, cooled to 0° and treated with HCl gas to give 26-3 as a tan solid.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.81 (m, 2H), 7.40 (m, 2H), 5.05 (t, J = 7.4 Hz, 1H), 3.72 (s, 3H), 3.30 (m, 2H).

3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (26-4)

The title compound 26-4 was prepared from 2-9A and 26-3 using the procedure described in Scheme 2. TLC R<sub>f</sub> = 0.40 (50% 20:1:1 EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O-50% EtOAc). FAB MS Obs'd 450.3 (M+H).

#### SCHEME 27



1-Methyl-4-bromopyrazole (27-2)

5 Methyl iodide (8.47 mL, 136 mmol) was added to a mixture of 4-bromopyrazole 27-1 (10 g, 38 mmol), and K<sub>2</sub>CO<sub>3</sub> (18.9 g, 136 mmol) in CH<sub>3</sub>CN (150 mL) and the mixture stirred at room temperature for 16 h, then filtered and evaporated to yield 27-2 as a yellow oil.  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.44(s, 1H), 7.38 (s, 1H), 3.90 (s, 3H).

10 3 (S)-Amino-3-(1-methyl-1H-pyrazol-4-yl)-propionic acid ethyl ester (27-3)

The bromide 27-2 was converted to the amino ester 27-3 following the procedure depicted in Scheme 1.  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.81 (s, 1H), 7.58 (s, 1H), 4.80 (m, 1H), 4.05 (q, J= 7.0 Hz, 2 H), 3.89 (s, 3H), 3.00 (m, 2 H), 1.24 (t, J = 7.0 Hz, 3 H).

15

3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (27-4)


The title compound 27-4 was prepared from 2-9A and 27-3 using the procedure described in Scheme 2.  
20 TLC R<sub>f</sub> = 0.24 (15:10:1:1 ethyl acetate/EtOH/water/NH<sub>4</sub>OH).  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.58 (s, 1H), 7.52 (d, J=7.3 Hz, 1H), 7.38 (s, 1H), 6.62 (d, J=7.3 Hz, 1H), 5.38 (m, 1H), 3.83 (s, 3H), 3.14-3.53 (9H), 2.97 (m, 2H), 2.80 (t, J=6.1 Hz, 2H), 2.67 (t, J=7.3 Hz, 2H), 1.93 (m, 4H).


25


30

35


**SCHEME A**  
**SYNTHESIS OF RADIOLIGAND FOR SPA ASSAY**



SCHEME A (CONTINUED)

SCHEME A. cont'd





**N-(4-Iodo-phenylsulfonylamino)-L-asparagine (A-2)**

To a stirred solution of acid A-1 (4.39 g, 33.2 mmol), NaOH (1.49 g, 37.2 mmol), dioxane (30 ml) and H<sub>2</sub>O (30 ml) at 0°C was added

5 pipsyl chloride (10.34 g, 34.2 mmol). After ~5 minutes, NaOH (1.49, 37.2 mmol), dissolved in 15 ml H<sub>2</sub>O, was added followed by the removal of the cooling bath. After 2.0 h, the reaction mixture was concentrated. The residue was dissolved in H<sub>2</sub>O (300 ml) and then washed with EtOAc. The aqueous portion was cooled to 0°C and then acidified with

10 concentrated HCl. The solid was collected and then washed with Et<sub>2</sub>O to provide acid A-2 as a white solid.

<sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ 7.86 (d, 2H, J=8Hz), 7.48 (d, 2H, J=8Hz) 3.70 (m, 1H), 2.39 (m, 2H).

15 **2(S)-(4-Iodo-phenylsulfonylamino)-β-alanine (A-3)**

To a stirred solution of NaOH (7.14 g, 181.8 mmol) and H<sub>2</sub>O (40 ml) at 0°C was added Br<sub>2</sub> (1.30 ml, 24.9 mmol) dropwise over a ten minute period. After ~5 minutes, acid A-2 (9.9 g, 24.9 mmol), NaOH (2.00 g, 49.8 mmol) and H<sub>2</sub>O (35 ml) were combined, cooled to 0°C and

20 then added in a single portion to the reaction. After stirring for

20 minutes at 0°C, the reaction was heated to 90°C for 30 minutes and then recooled to 0°C. The pH was adjusted to ~7 by dropwise addition of concentrated HCl. The solid was collected, washed with EtOAc, and then dried *in vacuo* to provide acid A-3 as a white solid.

5 <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ 8.02 (d, 2H, J=8Hz), 7.63 (d, 2H, J=8Hz), 4.36 (m, 1H), 3.51 (dd, 1H, J=5Hz, 13Hz) 3.21 (m, 1H).

Ethyl 2(S)-(4-iodo-phenylsulfonylamino)-β-alanine-hydrochloride (A-4)

HCl gas was rapidly bubbled through a suspension of acid  
10 A-3 (4.0 g, 10.81 mmol) in EtOH (50 ml) at 0°C for 10 minutes. The cooling bath was removed and the reaction was heated to 60°C. After 18 h, the reaction was concentrated to provide ester A-4 as a white solid.  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.98 (d, 2H, J=8Hz), 7.63 (d, 2H, J=8Hz),  
4.25 (q, 1H, J=5Hz), 3.92 (m, 2H), 3.33 (m, 1H), 3.06 (m, 1H), 1.01 (t, 3H,  
15 J=7Hz).

Ethyl 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoate (A-5a)

A mixture of ester A-5 (700 mg, 2.63 mmol), (for preparation, see: Scheme 29 of PCT International Application  
20 Publication No. WO 95/32710, published December 7, 1995) 10% Pd/C (350 mg) and EtOH were stirred under 1 atm H<sub>2</sub>. After 20 h, the reaction was filtered through a celite pad and then concentrated to provide ester A-5a as a brown oil.

TLC R<sub>f</sub> = 0.23 (silica, 40% EtOAc/hexanes)

25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.95 (d, 2H, J=8Hz), 7.26 (m, 3H), 6.43 (d, 1H, J=7Hz), 6.35 (d, 1H, J=8Hz), 4.37 (m, 4H), 3.05 (m, 2H), 2.91 (m, 2H), 1.39 (t, 3H, J=7Hz).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoic acid hydrochloride (A-6)

30 A suspension of ester A-5a (625 mg, 2.31 mmol) in 6N HCl (12 ml) was heated to 60°C. After ~20 h, the reaction was concentrated to give acid A-6 as a tan solid.

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.96 (d, 2H, J=8Hz), 7.80 (m, 1H), 7.33 (d, 2H, J=8Hz), 6.84 (d, 1H, J=9Hz), 6.69 (d, 1H, J=7Hz), 3.09 (m, 4H).

35

Ethyl 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-iodo-phenylsulfonylamino)- $\beta$ -alanine (A-7)

- A solution of acid 15-6 (400 mg, 1.43 mmol), amine A-4 (686 mg, 1.57 mmol), EDC (358 mg, 1.86 mmol), HOBT (252 mg, 1.86 mmol), NMM (632  $\mu$ l, 5.72 mmol) in DMF (10 ml) was stirred for ~20 h. The reaction was diluted with EtOAc and then washed with sat. NaHCO<sub>3</sub>, brine, dried (MgSO<sub>4</sub>) and concentrated. Flash chromatography (silica, EtOAc then 5% isopropanol/EtOAc) provided amide A-7 as a white solid.
- 10 TLC R<sub>f</sub> = 0.4 (silica, 10% isopropanol/EtOAc)  
<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD)  $\delta$  7.79 (d, 2H, J=9Hz) 7.61 (d, 2H, J=8Hz), 7.52 (d, 2H, J=9Hz), 7.29 (m, 1H), 7.27 (d, 2H, J=8Hz), 4.20 (m, 1H), 3.95 (q, 2H, J=7Hz), 3.66 (dd, 1H, J=6Hz, 14Hz), 3.49 (dd, 1H, J=8Hz, 13Hz), 3.01 (m, 2H), 2.86 (m, 2H), 1.08 (t, 3H, J=7Hz).

15 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-iodophenyl-sulfonylamino)- $\beta$ -alanine (A-8)

- A solution of ester A-7 (200 mg, 0.3213 mmol) and 6N HCl (30 ml) was heated to 60°C. After ~20 h, the reaction mixture was concentrated. Flash chromatography (silica, 20:20:1:1 EtOAc/EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O) provided acid A-8 as a white solid.
- 20 TLC R<sub>f</sub> = 0.45 (silica, 20:20:1:1 EtOAc/EtOH/NH<sub>4</sub>OH/H<sub>2</sub>O)  
<sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.40 (m, 1H), 8.14 (Bs, 1H), 7.81 (d, 2H, J=8Hz), 7.62 (d, 2H, J=8Hz), 7.48 (d, 2H, J=8Hz), 7.27 (m, 3H), 6.34 (d, 1H, J=7Hz), 6.25 (d, 1H, J=8Hz), 5.85 (bs, 2H), 3.89 (bs, 1H), 3.35 (m, 2H), 2.97 (m, 2H), 2.79 (m, 2H).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-trimethylstannyl-phenylsulfonylamino)- $\beta$ -alanine (A-9)

- 30 A solution of iodide A-8 (70 mg, 0.1178 mmol), [(CH<sub>3</sub>)<sub>3</sub>Sn]<sub>2</sub> (49  $\mu$ l, 0.2356 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mg) and dioxane (7 ml) was heated to 90°C. After 2 h, the reaction was concentrated and then purified by preparative HPLC (Delta-Pak C<sub>18</sub> 15  $\mu$ M 100A°, 40 x 100 mm; 95:5 then 5:95 H<sub>2</sub>O/CH<sub>3</sub>CN) to provide the trifluoroacetate salt. The salt was

suspended in H<sub>2</sub>O (10 ml), treated with NH<sub>4</sub>OH (5 drops) and then lyophilized to provide amide A-9 as a white solid.

<sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.40 (m, 1H), 8.18 (d, 1H, J=8Hz), 7.67 (m, 5H), 7.56 (d, 2H, J=8Hz), 7.29 (d, 2H, J=8Hz), 6.95-7.52 (m, 2H), 6.45 (bs, 2H), 4.00 (m, 1H), 3.50 (m, 1H), 3.33 (m, 1H), 2.97 (m, 2H), 2.86 (m, 2H).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-4-<sup>125</sup>iodo-phenylsulfonlamino-β-alanine (A-10)

10 An iodobead (Pierce) was added to a shipping vial of 5 mCi of Na<sup>125</sup>I (Amersham, IMS30) and stirred for five minutes at room temperature. A solution of 0.1 mg of A-9 in 0.05 mL of 10% H<sub>2</sub>SO<sub>4</sub>/MeOH was made and immediately added to the Na<sup>125</sup>I/iodobead vial. After stirring for three minutes at room temperature, approximately 0.04-0.05  
15 mL of NH<sub>4</sub>OH was added so the reaction mixture was at pH 6-7. The entire reaction mixture was injected onto the HPLC for purification [Vydac peptide-protein C-18 column, 4.6 x 250 mm, linear gradient of 10% acetonitrile (0.1% (TFA):H<sub>2</sub>O (0.1% TFA) to 90% acetonitrile (0.1% TFA):H<sub>2</sub>O (0.1% TFA) over 30 minutes, 1 mL/min]. The retention time  
20 of A-10 is 17 minutes under these conditions. Fractions containing the majority of the radioactivity were pooled, lyophilized and diluted with ethanol to give approximately 1 mCi of A-10, which coeluted on HPLC analysis with an authentic sample of A-8.

25 Instrumentation: Analytical and preparative HPLC was carried out using a Waters 600E Powerline Multi Solvent Delivery System with 0.1 mL heads with a Rheodyne 7125 injector and a Waters 990 Photodiode Array Detector with a Gilson FC203 Microfraction collector. For analytical and preparative HPLC, a Vydac peptide-protein  
30 C-18 column, 4.6 x 250 mm was used with a C-18 Brownlee modular guard column. The acetonitrile used for the HPLC analyses was Fisher Optima grade. The HPLC radiodetector used was a Beckman 170 Radioisotope detector. A Vydac C-18 protein and peptide column, 3.9 x 250 mm was used for analytical and preparative HPLC. Solutions of

radioactivity were concentrated using a Speedvac vacuum centrifuge. Calibration curves and chemical concentrations were determined using a Hewlett Packard Model 8452A UV/Vis Diode Array Spectrophotometer. Sample radioactivities were determined in a Packard A5530 gamma  
5 counter.

The test procedures employed to measure  $\alpha\nu\beta 3$  and  $\alpha\nu\beta 5$  binding and the bone resorption inhibiting activity of the compounds of the present invention are described below.

## 10 BONE RESORPTION-PIT ASSAY

When osteoclasts engage in bone resorption, they can cause the formation of pits in the surface of bone that they are acting upon. Therefore, when testing compounds for their ability to inhibit osteoclasts, it is useful to measure the ability of osteoclasts to excavate  
15 these resorption pits when the inhibiting compound is present.

Consecutive 200 micron thick cross sections from a 6 mm cylinder of bovine femur diaphysis are cut with a low speed diamond saw (Isomet, Beuler, Ltd., Lake Bluff, IL). Bone slices are pooled, placed in a 10% ethanol solution and refrigerated until further use.

20 Prior to experimentation, bovine bone slices are ultrasonicated twice, 20 minutes each in H<sub>2</sub>O. Cleaned slices are placed in 96 well plates such that two control lanes and one lane for each drug dosage are available. Each lane represents either triplicate or quadruplicate cultures. The bone slices in 96 well plates are sterilized by  
25 UV irradiation. Prior to incubation with osteoclasts, the bone slices are hydrated by the addition of 0.1 ml  $\alpha$ MEM, pH 6.9 containing 5% fetal bovine serum and 1% penicillin/streptomycin.

Long bones from 7-14 day old rabbits (New Zealand White Hare) are dissected, cleaned of soft tissue and placed in  $\alpha$ MEM  
30 containing 20 mM HEPES. The bones are minced using scissors until the pieces are <1 mm and transferred to a 50 ml tube in a volume of 25 ml. The tube is rocked gently by hand for 60 cycles, the tissue is sedimented for 1 min., and the supernatant is removed. Another 25 ml of medium is added to the tissue and rocked again. The second  
35 supernatant is combined with the first. The number of cells is counted

excluding erythrocytes (typically  $\sim 2 \times 10^7$  cells/ml). A cell suspension consisting of  $5 \times 10^6$ /ml in  $\alpha$ MEM containing 5% fetal bovine serum, 10 nM 1,25(OH) $_2$ D $_3$ , and pencillin-streptomycin is prepared. 200 ml aliquots are added to bovine bone slices (200 mm x 6 mm) and incubated  
5 for 2 hrs. at 37°C in a humidified 5% CO $_2$  atmosphere. The medium is removed gently with a micropipettor and fresh medium containing test compounds is added. The cultures are incubated for 48 hrs., and assayed for c-telopeptide (fragments of the  $\alpha$ 1 chain of type I collagen) by Crosslaps for culture media (Herlev, Denmark).

10 Bovine bone slices are exposed to osteoclasts for 20-24 hrs and are processed for staining. Tissue culture media is removed from each bone slice. Each well is washed with 200 ml of H $_2$ O, and the bone slices are then fixed for 20 minutes in 2.5% glutaraldehyde, 0.1 M cacodylate, pH 7.4. After fixation, any remaining cellular debris is  
15 removed by 2 min. ultrasonication in the presence of 0.25 M NH $_4$ OH followed by 2 X 15 min ultrasonication in H $_2$ O. The bone slices are immediately stained for 6-8 min with filtered 1% toluidine blue and 1% borax.

After the bone slices have dried, resorption pits are counted  
20 in test and control slices. Resorption pits are viewed in a Microphot Fx (Nikon) fluorescence microscope using a polarizing Nikon IGS filter cube. Test dosage results are compared with controls and resulting IC $_{50}$  values are determined for each compound tested.

The appropriateness of extrapolating data from this assay to  
25 mammalian (including human) disease states is supported by the teaching found in Sato, M., et al., Journal of Bone and Mineral Research, Vol. 5, No. 1, pp.31-40, 1990, which is incorporated by reference herein in its entirety. This article teaches that certain bisphosphonates have been used clinically and appear to be effective in  
30 the treatment of Paget's disease, hypercalcemia of malignancy, osteolytic lesions produced by bone metastases, and bone loss due to immobilization or sex hormone deficiency. These same bisphosphonates are then tested in the resorption pit assay described above to confirm a

correlation between their known utility and positive performance in the assay.

#### EIB ASSAY

- 5                   Duong *et al.*, *J. Bone Miner. Res.*, 8: S378 (1993) describes a system for expressing the human integrin  $\alpha\beta 3$ . It has been suggested that the integrin stimulates attachment of osteoclasts to bone matrix, since antibodies against the integrin, or RGD-containing molecules, such as echistatin (European Publication 382 451), can effectively block
- 10 bone resorption.

#### Reaction Mixture:

1. 175  $\mu$ l TBS buffer (50 mM Tris•HCl pH 7.2, 150 mM NaCl, 1% BSA, 1 mM  $\text{CaCl}_2$ , 1 mM  $\text{MgCl}_2$ ).
- 15 2. 25  $\mu$ l cell extract (dilute with 100 mM octylglucoside buffer to give 2000 cpm/25  $\mu$ l).
3.  $^{125}\text{I}$ -echistatin (25  $\mu$ l/50,000 cpm) (see EP 382 451).
4. 25  $\mu$ l buffer (total binding) or unlabeled echistatin (non-specific binding).

20

The reaction mixture was then incubated for 1 h at room temp. The unbound and the bound  $\alpha\beta 3$  were separated by filtration using a Skatron Cell Harvester. The filters (prewet in 1.5% polyethyleneimine for 10 mins) were then washed with the wash buffer (50

25 mM Tris HCl, 1mM  $\text{CaCl}_2/\text{MgCl}_2$ , pH 7.2). The filter was then counted in a gamma counter.

#### SPA ASSAY

#### 30 MATERIALS:

1. Wheat germ agglutinin Scintillation Proximity Beads (SPA): Amersham
2. Octylglucopyranoside: Calbiochem
- 35 3. HEPES: Calbiochem

4. NaCl: Fisher
5. CaCl<sub>2</sub>: Fisher
6. MgCl<sub>2</sub>: SIGMA
7. Phenylmethylsulfonylfluoride (PMSF): SIGMA
- 5 8. Optiplate: PACKARD
9. Compound A-10 (specific activity 500-1000 Ci/mmol)
10. test compound
11. Purified integrin receptor:  $\alpha_v\beta_3$  was purified from 293 cells overexpressing  $\alpha_v\beta_3$  (Duong *et al.*, *J. Bone Min. Res.*, 8:S378, 1993) according to Pytela (*Methods in Enzymology*, 144:475, 1987)
12. Binding buffer: 50 mM HEPES, pH 7.8, 100 mM NaCl, 1 mM Ca<sup>2+</sup>/Mg<sup>2+</sup>, 0.5 mM PMSF
13. 50 mM octylglucoside in binding buffer: 50-OG buffer

15

## PROCEDURE:

1. Pretreatment of SPA beads:  
20 500 mg of lyophilized SPA beads were first washed four times with 200 ml of 50-OG buffer and once with 100 ml of binding buffer, and then resuspended in 12.5 ml of binding buffer.
2. Preparation of SPA beads and receptor mixture  
25 In each assay tube, 2.5  $\mu$ l (40 mg/ml) of pretreated beads were suspended in 97.5  $\mu$ l of binding buffer and 20  $\mu$ l of 50-OG buffer. 5  $\mu$ l (~30 ng/ $\mu$ l) of purified receptor was added to the beads in suspension with stirring at room temperature for 30 minutes. The mixture was then centrifuged at 2,500 rpm in a Beckman GPR Benchtop centrifuge for 10 minutes at 4°C. The  
30 pellets were then resuspended in 50  $\mu$ l of binding buffer and 25  $\mu$ l of 50-OG buffer.



3. Reaction

The following were sequentially added into Optiplate in corresponding wells:

- 5 (i) Receptor/beads mixture (75  $\mu$ l)  
(ii) 25  $\mu$ l of each of the following: compound to be tested, binding buffer for total binding or A-8 for non-specific binding (final concentration 1  $\mu$ M)  
10 (iii) A-10 in binding buffer (25  $\mu$ l, final concentration 40 pM)  
(iv) Binding buffer (125  $\mu$ l)  
(v) Each plate was sealed with plate sealer from PACKARD and incubated overnight with rocking at 4°C

15 4. Plates were counted using PACKARD TOPCOUNT

5. % inhibition was calculated as follows:

A = total counts

B = nonspecific counts

20 C = sample counts

% inhibition =  $[(A-B)-(C-B)]/(A-B)/(A-B) \times 100$

## OCFORM ASSAY

Osteoblast-like cells (1.8 cells), originally derived from mouse calvaria, were plated in CORNING 24 well tissue culture plates in  $\alpha$ MEM medium containing ribo- and deoxyribonucleosides, 10% fetal bovine serum and penicillin-streptomycin. Cells were seeded at 40,000/well in the morning. In the afternoon, bone marrow cells were prepared from six week old male Balb/C mice as follows:

30 Mice were sacrificed, tibiae removed and placed in the above medium. The ends were cut off and the marrow was flushed out of the cavity into a tube with a 1 mL syringe with a 27.5 gauge needle. The marrow was suspended by pipetting up and down. The suspension was passed through >100  $\mu$ m nylon cell strainer. The resulting suspension was centrifuged at 350 x g for seven minutes. The pellet was

resuspended, and a sample was diluted in 2% acetic acid to lyse the red cells. The remaining cells were counted in a hemacytometer. The cells were pelleted and resuspended at  $1 \times 10^6$  cells/mL. 50  $\mu$ L was added to each well of 1.8 cells to yield 50,000 cells/well and 1,25-dihydroxy-vitamin D3 (D3) was added to each well to a final concentration of 10 nM. The cultures were incubated at 37°C in a humidified, 5% CO<sub>2</sub> atmosphere. After 48 h, the medium was changed. 72 h after the addition of bone marrow, test compounds were added with fresh medium containing D3 to quadruplicate wells. Compounds were added again after 48 h with fresh medium containing D3. After an additional 48 h., the medium was removed, cells were fixed with 10% formaldehyde in phosphate-buffered saline for 10 minutes at room temperature, followed by a 1-2 minute treatment with ethanol:acetone (1:1) and air dried. The cells were then stained for tartrate resistant acid phosphatase as follows:

The cells were stained for 10-15 minutes at room temperature with 50 mM acetate buffer, pH 5.0 containing 30 mM sodium tartrate, 0.3 mg/mL Fast Red Violet LB Salt and 0.1 mg/mL Naphthol AS -MX phosphate. After staining, the plates were washed extensively with deionized water and air dried. The number of multinucleated, positive staining cells was counted in each well.

#### $\alpha$ v $\beta$ 5 ATTACHMENT ASSAY

Duong *et al.*, *J. Bone Miner. Res.*, 11: S290 (1996), describes a system for expressing the human  $\alpha$ v $\beta$ 5 integrin receptor.

#### Materials:

1. Media and solutions used in this assay are purchased from BRL/Gibco, except BSA and the chemicals are from Sigma.
2. Attachment medium: HBSS with 1 mg/ml heat-inactivated fatty acid free BSA and 2 mM CaCl<sub>2</sub>.
3. Glucosaminidase substrate solution: 3.75 mM p-nitrophenyl N-acetyl-beta-D-glucosaminide, 0.1 M sodium citrate, 0.25% Triton, pH 5.0.

4. Glycine-EDTA developing solution: 50 mM glycine, 5 mM EDTA, pH 10.5.

Methods:

- 5           1. Plates (96 well, Nunc Maxi Sorp) were coated overnight at 4°C with human vitronectin (3 ug/ml) in 50 mM carbonate buffer (pH 9.6), using 100 µl/well. Plates were then washed 2X with DPBS and blocked with 2% BSA in DPBS for 2h at room temperature. After additional washes (2X) with DPBS, plates were used for cell attachment assay.
- 10           2. 293 (αvβ5) cells were grown in MEM media in presence of 10% fetal calf serum to 90% confluence. Cells were then lifted from dishes with 1X Trypsin/EDTA and washed 3X with serum free MEM. Cells were resuspended in attachment medium (3 X 10<sup>5</sup> cells/ml).
- 15           3. Test compounds were prepared as a series of dilutions at 2X concentrations and added as 50 µl/well. Cell suspension was then added as 50 µl/well. Plates were incubated at 37°C with 55 CO<sub>2</sub> for 1 hour to allow attachment.
- 20           4. Non-adherent cells were removed by gently washing the plates (3X) with DPBS and then incubated with glucosaminidase substrate solution (100 µl/well), overnight at room temperature in the dark. To quantitate cell numbers, standard curve of glucosaminidase activity was determined for each experiment by adding samples of cell suspension directly to wells containing the enzyme substrate solution.
- 25           5. The next day, the reaction was developed by addition of 185 µl/well of glycine/EDTA solution and reading absorbance at 405 nm using a Molecular Devices V-Max plate reader. Average test absorbance values (4 wells per test samples) were calculated. Then, the number of attached cells at each drug concentration was quantitated versus the standard curve of cells using the Softmax program.
- 30
- 35

### EXAMPLE OF A PHARMACEUTICAL FORMULATION

As a specific embodiment of an oral composition, 100 mg of a compound of the present invention are formulated with sufficient  
5 finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.





Representative compounds of the present invention were tested and found to bind to human  $\alpha v \beta 3$  integrin. These compounds are generally found to have IC<sub>50</sub> values less than about 100 nM in the SPA  
10 assay.



Representative compounds of the present invention were tested and generally found to inhibit  $\geq 50\%$  the attachment of  $\alpha v \beta 5$  expressing cells to plates coated with vitronectin at concentrations of about 1  $\mu$ M.

15 While the invention has been described and illustrated in reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the preferred doses  
20 as set forth hereinabove may be applicable as a consequence of variations in the responsiveness of the mammal being treated for severity of bone disorders caused by resorption, or for other indications for the compounds of the invention indicated above. Likewise, the specific pharmacological responses observed may vary according to and  
25 depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended,  
30 therefore, that the invention be limited only by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.


## WHAT IS CLAIMED IS:

1. A compound having a structural formula selected from the group consisting of





, and




5

;

wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are substituted only with R<sup>10</sup> and R<sup>12</sup>;

10

X is selected from the group consisting of



15

a 5- or 6-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R<sup>1</sup> substituent and the ring

carbon atoms are unsubstituted or substituted with one or two  $R^1$  substituents, and

5 a 9- to 14-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one  $R^1$  substituent and the ring carbon atoms are unsubstituted or substituted with one or two  $R^1$  substituents;

10

Y is selected from the group consisting of

$-(CH_2)_m-$ ,  
 $-(CH_2)_m-O-(CH_2)_n-$ ,  
 $-(CH_2)_m-NR^4-(CH_2)_n-$ ,  
15  $-(CH_2)_m-S-(CH_2)_n-$ ,  
 $-(CH_2)_m-SO-(CH_2)_n-$ ,  
 $-(CH_2)_m-SO_2-(CH_2)_n-$ ,  
 $-(CH_2)_m-O-(CH_2)_n-O-(CH_2)_p-$ ,  
 $-(CH_2)_m-O-(CH_2)_n-NR^4-(CH_2)_p-$ ,  
20  $-(CH_2)_m-NR^4-(CH_2)_n-NR^4-(CH_2)_p-$ ,  
 $-(CH_2)_m-O-(CH_2)_n-S-(CH_2)_p-$ ,  
 $-(CH_2)_m-S-(CH_2)_n-S-(CH_2)_p-$ ,  
 $-(CH_2)_m-NR^4-(CH_2)_n-S-(CH_2)_p-$ ,  
 $-(CH_2)_m-NR^4-(CH_2)_n-O-(CH_2)_p-$ ,  
25  $-(CH_2)_m-S-(CH_2)_n-O-(CH_2)_p-$ ,  
 $-(CH_2)_m-S-(CH_2)_n-NR^4-(CH_2)_p-$ , and  
 $-(CH_2)_m-Z-(CH_2)_n-$ ,

30 wherein Z is a 3- to 10-membered monocyclic or polycyclic aromatic or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one  $R^1$  substituent and the ring carbon atoms are unsubstituted or substituted with one or two  $R^1$  substituents, and

wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than in R<sup>4</sup>, can be substituted by one or two R<sup>3</sup> substituents; and

- wherein R<sup>1</sup> and R<sup>2</sup> are each independently selected from the group
- 5 consisting of
- hydrogen, halogen, C<sub>1-10</sub> alkyl, C<sub>3-8</sub> cycloalkyl,  
 C<sub>3-8</sub> cycloheteroalkyl, C<sub>3-8</sub> cycloalkyl C<sub>1-6</sub> alkyl,  
 C<sub>3-8</sub> cycloheteroalkyl C<sub>1-6</sub> alkyl, aryl, aryl C<sub>1-8</sub> alkyl, amino,  
 amino C<sub>1-8</sub> alkyl, C<sub>1-3</sub> acylamino, C<sub>1-3</sub> acylamino C<sub>1-8</sub> alkyl,  
 10 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino, (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-8</sub> alkyl,  
 C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkoxy C<sub>1-6</sub> alkyl, hydroxycarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl, C<sub>1-3</sub> alkoxycarbonyl,  
 C<sub>1-3</sub> alkoxycarbonyl C<sub>1-6</sub> alkyl, hydroxycarbonyl-  
 C<sub>1-6</sub> alkyloxy, hydroxy, hydroxy C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkyloxy-  
 15 C<sub>1-6</sub> alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,  
 trifluoroethoxy, C<sub>1-8</sub> alkyl-S(O)<sub>p</sub>, (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-8</sub> alkyloxycarbonylamino, (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>amino, (aryl)<sub>p</sub>amino, aryl C<sub>1-8</sub>  
 alkylsulfonylamino, and C<sub>1-8</sub> alkylsulfonylamino;  
 20 or two R<sup>1</sup> substituents, when on the same carbon atom, are taken  
 together with the carbon atom to which they are attached to  
 form a carbonyl group;

- each R<sup>3</sup> is independently selected from the group consisting of
- 25 hydrogen,  
 aryl,  
 C<sub>1-10</sub> alkyl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 30 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 halogen,



- hydroxyl,  
oxo,  
trifluoromethyl,  
C<sub>1-8</sub> alkylcarbonylamino,  
5 aryl C<sub>1-5</sub> alkoxy,  
C<sub>1-5</sub> alkoxycarbonyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
C<sub>1-6</sub> alkylcarbonyloxy,  
C<sub>3-8</sub> cycloalkyl,  
10 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
amino C<sub>1-6</sub> alkyl,  
arylamino carbonyl,  
aryl C<sub>1-5</sub>-alkylaminocarbonyl,  
aminocarbonyl,  
15 aminocarbonyl C<sub>1-6</sub> alkyl,  
hydroxycarbonyl,  
hydroxycarbonyl C<sub>1-6</sub> alkyl,  
HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
20 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
25 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
30 C<sub>1-6</sub> alkoxy,  
aryl C<sub>1-6</sub> alkoxy,  
aryl C<sub>1-6</sub> alkyl,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>amino,  
35 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,

- (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 5 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 10 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkoxycarbonylamino,  
 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryloxy carbonylamino C<sub>1-8</sub> alkyl,  
 15 aryl C<sub>1-8</sub> alkoxycarbonylamino,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 arylcarbonylamino C<sub>1-6</sub> alkyl,  
 20 aryl C<sub>1-6</sub> alkylcarbonylamino,  
 aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 25 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 30 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylsulfonyl,

C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 arylsulfonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 5 C<sub>1-6</sub> alkylcarbonyl,  
 C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 arylcarbonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 10 C<sub>1-6</sub> alkylthiocarbonylamino,  
 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 15 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl;  
 or two R<sup>3</sup> substituents, when on the same carbon atom, are taken  
 20 together with the carbon atom to which they are attached to  
 form a carbonyl group or a cyclopropyl group,  
 wherein any of the alkyl groups of R<sup>3</sup> are either unsubstituted or  
 substituted with one to three R<sup>1</sup> substituents, provided that each R<sup>3</sup> is  
 selected such that in the resultant compound the carbon atom or atoms  
 25 to which R<sup>3</sup> is attached is itself attached to no more than one  
 heteroatom;

each R<sup>4</sup> is independently selected from the group consisting of  
 hydrogen,  
 30 aryl,  
 aminocarbonyl,  
 C<sub>3-8</sub> cycloalkyl,  
 amino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl,

- (aryl C<sub>1-5</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkyl,  
 aryl C<sub>1-6</sub> alkyl,  
 5 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
 C<sub>1-8</sub> alkylsulfonyl,  
 C<sub>1-8</sub> alkoxy carbonyl,  
 aryloxy carbonyl,  
 10 aryl C<sub>1-8</sub> alkoxy carbonyl,  
 C<sub>1-8</sub> alkyl carbonyl,  
 aryl carbonyl,  
 aryl C<sub>1-6</sub> alkyl carbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 15 aminosulfonyl,  
 C<sub>1-8</sub> alkyl aminosulfonyl,  
 (aryl)<sub>p</sub>aminosulfonyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonyl,  
 arylsulfonyl,  
 20 aryl C<sub>1-6</sub> alkylsulfonyl,  
 C<sub>1-6</sub> alkylthiocarbonyl,  
 arylthiocarbonyl, and  
 aryl C<sub>1-6</sub> alkylthiocarbonyl,

wherein any of the alkyl groups of R<sup>4</sup> are either unsubstituted or  
 25 substituted with one to three R<sup>1</sup> substituents;

R<sup>5</sup> and R<sup>6</sup> are each independently selected from the group consisting of  
 hydrogen,  
 C<sub>1-10</sub> alkyl,  
 30 aryl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,

- aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 halogen,  
 hydroxyl,  
 5 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 10 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 amino C<sub>1-6</sub> alkyl,  
 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 15 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 hydroxycarbonyl,  
 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 20 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 25 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 30 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,

- (aryl)pamino,  
(aryl)pamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-6</sub> alkyl)pamino,  
(aryl C<sub>1-6</sub> alkyl)pamino C<sub>1-6</sub> alkyl,  
5 arylcarbonyloxy,  
aryl C<sub>1-6</sub> alkylcarbonyloxy,  
(C<sub>1-6</sub> alkyl)paminocarbonyloxy,  
C<sub>1-8</sub> alkylsulfonylamino,  
arylsulfonylamino,  
10 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
arylsulfonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylsulfonylamino,  
aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-8</sub> alkoxycarbonylamino,  
15 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
aryl C<sub>1-8</sub> alkoxycarbonylamino,  
aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonylamino,  
20 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonylamino,  
aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
25 (C<sub>1-8</sub> alkyl)paminocarbonylamino,  
(C<sub>1-8</sub> alkyl)paminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl)paminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)paminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)paminocarbonylamino C<sub>1-6</sub> alkyl,  
30 aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)paminosulfonylamino,  
(C<sub>1-8</sub> alkyl)paminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl)paminosulfonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)paminosulfonylamino,

- (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylsulfonyl,  
 C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 arylsulfonyl C<sub>1-6</sub> alkyl,  
 5 aryl C<sub>1-6</sub> alkylsulfonyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylcarbonyl,  
 C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 arylcarbonyl C<sub>1-6</sub> alkyl,  
 10 aryl C<sub>1-6</sub> alkylcarbonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylthiocarbonylamino,  
 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 15 aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
 20 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl;  
 or R<sup>5</sup> and R<sup>6</sup> are taken together with the carbon atom to which they are  
 attached to form a carbonyl group,  
 wherein any of the alkyl groups of R<sup>5</sup> or R<sup>6</sup> are either unsubstituted or  
 substituted with one to three R<sup>1</sup> substituents,  
 25 and provided that each R<sup>5</sup> and R<sup>6</sup> are selected such that in the resultant  
 compound the carbon atom to which R<sup>5</sup> and R<sup>6</sup> are attached is itself  
 attached to no more than one heteroatom;  
  
 R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of  
 30 hydrogen,  
 C<sub>1-10</sub> alkyl,  
 aryl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,

- aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 5 halogen,  
 hydroxyl,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 10 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 C<sub>3-8</sub> cycloalkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 amino C<sub>1-6</sub> alkyl,  
 15 arylaminocarbonyl,  
 aryl C<sub>1-5</sub> alkylaminocarbonyl,  
 aminocarbonyl,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 hydroxycarbonyl,  
 20 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 25 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 30 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkoxy,  
 aryl C<sub>1-6</sub> alkoxy,



- aryl C<sub>1-6</sub> alkyl,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
(aryl)<sub>p</sub>amino,  
(aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
5 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
(aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
arylcarbonyloxy,  
aryl C<sub>1-6</sub> alkylcarbonyloxy,  
(C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
10 C<sub>1-8</sub> alkylsulfonylamino,  
arylcarbonylamino,  
arylsulfonylamino,  
C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
arylsulfonylamino C<sub>1-6</sub> alkyl,  
15 aryl C<sub>1-6</sub> alkylsulfonylamino,  
aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
C<sub>1-8</sub> alkoxycarbonylamino,  
C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
20 aryl C<sub>1-8</sub> alkoxycarbonylamino,  
aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
arylcarbonylamino C<sub>1-6</sub> alkyl,  
aryl C<sub>1-6</sub> alkylcarbonylamino,  
25 aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
aminocarbonylamino C<sub>1-6</sub> alkyl,  
arylaminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
30 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
(aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
aminosulfonylamino C<sub>1-6</sub> alkyl,  
(C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,

- (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 5 C<sub>1-6</sub> alkylsulfonyl,  
 C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 arylsulfonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 10 C<sub>1-6</sub> alkylcarbonyl,  
 C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 arylcarbonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 15 C<sub>1-6</sub> alkylthiocarbonylamino,  
 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 20 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl, and  
 C<sub>7-20</sub> polycyclyl C<sub>0-8</sub> alkylsulfonylamino;  
 25 wherein any of the alkyl groups of R<sup>7</sup> and R<sup>8</sup> are either unsubstituted or  
 substituted with one to three R<sup>1</sup> substituents,  
 and provided that each R<sup>7</sup> and R<sup>8</sup> are selected such that in the resultant  
 compound the carbon atom to which R<sup>7</sup> and R<sup>8</sup> are attached is itself  
 attached to no more than one heteroatom;  
 30 R<sup>9</sup> is selected from the group consisting of  
 hydrogen,  
 C<sub>1-8</sub> alkyl,  
 aryl,

- aryl C<sub>1-8</sub> alkyl,  
 C<sub>1-8</sub> alkylcarbonyloxy C<sub>1-4</sub> alkyl,  
 aryl C<sub>1-8</sub> alkylcarbonyloxy C<sub>1-4</sub> alkyl,  
 C<sub>1-8</sub> alkylaminocarbonylmethylene, and  
 5 C<sub>1-8</sub> dialkylaminocarbonylmethylene;

R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub> and R<sub>13</sub> are each independently selected from the group consisting of

- 10 hydrogen,  
 C<sub>1-8</sub> alkyl,  
 aryl,  
 halogen,  
 hydroxyl;  
 15 aminocarbonyl,  
 C<sub>3-8</sub> cycloalkyl,  
 amino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl,  
 hydroxycarbonyl,  
 (aryl C<sub>1-5</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 20 hydroxycarbonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkyl,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>2-6</sub> alkyl,  
 C<sub>1-8</sub> alkylsulfonyl,  
 25 C<sub>1-8</sub> alkoxycarbonyl,  
 aryloxy carbonyl,  
 aryl C<sub>1-8</sub> alkoxycarbonyl,  
 C<sub>1-8</sub> alkylcarbonyl,  
 arylcarbonyl,  
 30 aryl C<sub>1-6</sub> alkylcarbonyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 aminosulfonyl,  
 C<sub>1-8</sub> alkylaminosulfonyl,  
 (aryl)<sub>p</sub>aminosulfonyl,

- (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonyl,  
 C<sub>1-6</sub> alkylsulfonyl,  
 arylsulfonyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl,  
 5 aryl C<sub>1-6</sub> alkylcarbonyl,  
 C<sub>1-6</sub> alkylthiocarbonyl,  
 arylthiocarbonyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonyl,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-O-(CH<sub>2</sub>)<sub>s</sub>-,  
 10 aryl-(CH<sub>2</sub>)<sub>r</sub>S(O)<sub>p</sub>-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-C(O)-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-C(O)-(CH<sub>2</sub>)<sub>s</sub>-,  
 aryl-(CH<sub>2</sub>)<sub>r</sub>-N(R<sup>4</sup>)-(CH<sub>2</sub>)<sub>s</sub>-,  
 15 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
 20 CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>3-7</sub> cycloalkyl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 aryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-CH=CH-(CH<sub>2</sub>)<sub>t</sub>-,  
 25 C<sub>1-6</sub> alkyl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-6</sub> alkylaryl-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>t</sub>-,  
 C<sub>1-8</sub> alkylcarbonylamino,  
 aryl C<sub>1-5</sub> alkoxy,  
 C<sub>1-5</sub> alkoxycarbonyl,  
 30 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl,  
 C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 aminocarbonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkoxy,

- aryl C<sub>1-6</sub> alkoxy,  
 (aryl)<sub>p</sub>amino,  
 (aryl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino,  
 5 (aryl C<sub>1-6</sub> alkyl)<sub>p</sub>amino C<sub>1-6</sub> alkyl,  
 arylcarbonyloxy,  
 aryl C<sub>1-6</sub> alkylcarbonyloxy,  
 (C<sub>1-6</sub> alkyl)<sub>p</sub>aminocarbonyloxy,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 10 arylsulfonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 15 C<sub>1-8</sub> alkoxycarbonylamino,  
 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryloxy carbonylamino C<sub>1-8</sub> alkyl,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 20 C<sub>1-8</sub> alkylcarbonylamino,  
 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 arylcarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylcarbonylamino,  
 aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 25 aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 30 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,

- (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylsulfonyl,  
 C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 5 arylsulfonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl,  
 aryl C<sub>1-6</sub> alkylsulfonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylcarbonyl,  
 C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 10 arylcarbonyl C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl,  
 aryl C<sub>1-6</sub> alkylcarbonyl C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylthiocarbonylamino,  
 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 15 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl,  
 20 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl, and  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonyl C<sub>1-6</sub> alkyl; or  
 R<sup>10</sup> and R<sup>12</sup> are taken together with the carbon atoms to which  
 they are attached to form a 5- to 7-membered monocyclic aromatic  
 or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms  
 25 selected from the group consisting of N, O, and S wherein the ring  
 nitrogen atoms are unsubstituted or substituted with one R<sup>1</sup>  
 substituent and the ring carbon atoms are unsubstituted or  
 substituted with one or two R<sup>1</sup> substituents,  
 and wherein any of the alkyl groups of R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> are either  
 30 unsubstituted or substituted with one to three R<sup>1</sup> substituents;

wherein



each m is independently an integer from 0 to 6;

each n is independently an integer from 0 to 6


- each p is independently an integer from 0 to 2;  
 each r is independently an integer from 1 to 3;  
 each s is independently an integer from 0 to 3;  
 each t is independently an integer from 0 to 3; and  
 5 each v is independently an integer from 0 to 2;

and the pharmaceutically acceptable salts thereof.

2. The compound of Claim 1 having a structural  
 10 formula selected from the group consisting of



and



15

wherein the dotted line **a** represents a single or a double bond, provided that when **a** represents a double bond, the double bond carbon atoms are substituted only with R<sup>10</sup> and R<sup>12</sup>;


20

X is

a 6-membered monocyclic aromatic ring system having 1 or 2 nitrogen atoms wherein each ring carbon atom is unsubstituted or substituted with one R<sup>1</sup> substituent, or

- 5 a 9- to 14-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R<sup>1</sup> substituent and the ring carbon atoms are
- 10 unsubstituted or substituted with one or two R<sup>1</sup> substituents.

3. The compound of Claim 2 having structural formula




15

wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are


20 substituted only with R<sup>10</sup> and R<sup>12</sup>; and

X is selected from the group consisting of






4. The compound of Claim 3 having structural formula



5

5. The compound of Claim 4 wherein X is



6. The compound of Claim 5 wherein Y is selected from the group consisting of

10

-(CH<sub>2</sub>)<sub>m</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>-,

15

-(CH<sub>2</sub>)<sub>m</sub>-SO-(CH<sub>2</sub>)<sub>n</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-SO<sub>2</sub>-(CH<sub>2</sub>)<sub>n</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-O-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-,

-(CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>p</sub>-, and

20

-(CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>-O-(CH<sub>2</sub>)<sub>p</sub>-,

wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than in R<sup>4</sup>, can be substituted by one or two R<sup>3</sup> substituents.

25

7. The compound of Claim 6 wherein Y is selected from the group consisting of

(CH<sub>2</sub>)<sub>m</sub>, (CH<sub>2</sub>)<sub>m</sub>-S-(CH<sub>2</sub>)<sub>n</sub>, and (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>,  
5 wherein any methylene (CH<sub>2</sub>) carbon atom in Y, other than in R<sup>4</sup>, can be substituted by one or two R<sup>3</sup> substituents, m and n are integers from 0-4, and v is 0.

10 8. The compound of Claim 7 wherein Y is

(CH<sub>2</sub>)<sub>m</sub> or (CH<sub>2</sub>)<sub>m</sub>-NR<sup>4</sup>-(CH<sub>2</sub>)<sub>n</sub>,  
wherein any methylene (CH<sub>2</sub>) group in Y, other than in R<sup>4</sup>, can be substituted by one or two R<sup>3</sup> substituents.

15

9. The compound of Claim 8 wherein each R<sup>3</sup> is independently selected from the group consisting of

hydrogen,  
fluoro,  
20 trifluoromethyl,  
aryl,  
C<sub>1-8</sub> alkyl,  
arylC<sub>1-6</sub> alkyl  
hydroxyl,  
25 oxo,  
arylaminocarbonyl,  
aryl C<sub>1-5</sub> alkylaminocarbonyl,  
aminocarbonyl, and  
aminocarbonyl C<sub>1-6</sub> alkyl;

30

and each R<sup>4</sup> is independently selected from the group consisting of

hydrogen,  
aryl,  
C<sub>3-8</sub> cycloalkyl,  
35 C<sub>1-8</sub> alkyl,

C<sub>1-8</sub> alkylcarbonyl,  
arylcarbonyl,  
C<sub>1-6</sub> alkylsulfonyl,  
arylsulfonyl,  
5 arylC<sub>1-6</sub>alkylsulfonyl,  
arylC<sub>1-6</sub>alkylcarbonyl,  
C<sub>1-8</sub>alkylaminocarbonyl,  
arylC<sub>1-5</sub>alkylaminocarbonyl,  
arylC<sub>1-8</sub>alkoxycarbonyl, and  
10 C<sub>1-8</sub>alkoxycarbonyl.

10. The compound of Claim 9 wherein R<sup>6</sup>, R<sup>7</sup>, and R<sup>8</sup> are each hydrogen and R<sup>5</sup> is selected from the group consisting of  
hydrogen,  
15 aryl,  
C<sub>1-8</sub> alkyl,  
aryl-C≡C-(CH<sub>2</sub>)<sub>t</sub>-,  
aryl C<sub>1-6</sub> alkyl,  
CH<sub>2</sub>=CH-(CH<sub>2</sub>)<sub>t</sub>-, and  
20 HC≡C-(CH<sub>2</sub>)<sub>t</sub>-.

11. The compound of Claim 10 wherein R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> are each independently selected from the group consisting of  
hydrogen, aryl, C<sub>1-6</sub>alkyl, and arylC<sub>1-6</sub>alkyl.

25

12. The compound of Claim 10 wherein R<sup>9</sup> is selected from the group consisting of hydrogen, methyl, and ethyl.

13. The compound of Claim 12 wherein R<sup>9</sup> is hydrogen.

30

14. The compound of Claim 9 wherein R<sup>5</sup>, R<sup>6</sup>, and R<sup>8</sup> are each hydrogen and R<sup>7</sup> is selected from the group consisting of  
hydrogen,  
aryl,

- C<sub>1-8</sub> alkylcarbonylamino,  
 C<sub>1-8</sub> alkylsulfonylamino,  
 arylcarbonylamino,  
 arylsulfonylamino,  
 5 C<sub>1-8</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 arylsulfonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylsulfonylamino,  
 aryl C<sub>1-6</sub> alkylsulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-8</sub> alkoxycarbonylamino,  
 10 C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryloxycarbonylamino C<sub>1-8</sub> alkyl,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino,  
 aryl C<sub>1-8</sub> alkoxycarbonylamino C<sub>1-8</sub> alkyl,  
 C<sub>1-8</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 15 arylcarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylcarbonylamino,  
 aryl C<sub>1-6</sub> alkylcarbonylamino C<sub>1-6</sub> alkyl,  
 aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 20 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 arylaminocarbonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino C<sub>1-6</sub> alkyl,  
 25 aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino,  
 30 (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino C<sub>1-6</sub> alkyl,  
 C<sub>1-6</sub> alkylthiocarbonylamino,  
 C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 arylthiocarbonylamino C<sub>1-6</sub> alkyl,  
 aryl C<sub>1-6</sub> alkylthiocarbonylamino, and

aryl C<sub>1-6</sub> alkylthiocarbonylamino C<sub>1-6</sub> alkyl.

15. The compound of Claim 14 wherein R<sup>7</sup> is selected from the group consisting of

- 5        hydrogen,  
         aryl,  
         C<sub>1-8</sub> alkylcarbonylamino,  
         aryl C<sub>1-6</sub> alkylcarbonylamino,  
         arylcarbonylamino,  
10       C<sub>1-8</sub> alkylsulfonylamino,  
         aryl C<sub>1-6</sub> alkylsulfonylamino,  
         arylsulfonylamino,  
         C<sub>1-8</sub> alkoxy carbonylamino,  
         aryl C<sub>1-8</sub> alkoxy carbonylamino,  
15       arylaminocarbonylamino,  
         (C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
         (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminocarbonylamino,  
         (C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino, and  
         (aryl C<sub>1-8</sub> alkyl)<sub>p</sub>aminosulfonylamino.

20

16. The compound of Claim 15 wherein R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> are each independently selected from the group consisting of hydrogen, aryl, C<sub>1-6</sub>alkyl, and arylC<sub>1-6</sub> alkyl.

25

17. The compound of Claim 15 wherein R<sup>9</sup> is selected from the group consisting of hydrogen, methyl, and ethyl.

30

18. The compound of Claim 17 wherein R<sup>9</sup> is hydrogen.

19. The compound of Claim 9 selected from the group consisting of

- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(R)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35 3-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 5 3(R)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 10 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,
- 3(R)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,
- 15 3-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,
- 20 3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 30 3(R)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,  
trifluoroacetate salt,
- 5 3(S)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,  
bis(trifluoroacetate) salt,
- 10 3(R)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 15 3(S)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 20 3-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-  
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-  
propionic acid,
- 30 3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-  
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-  
propionic acid,
- 35 3-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-  
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-  
propionic acid,



- 3(S)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
5 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
20 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
35 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 2(S)-Benzenesulfonylamino-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-  
imidazolidin-1-yl)-pent-4-enoic acid,
- 25 3(R)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-  
imidazolidin-1-yl)-pent-4-enoic acid,
- 3-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-  
30 imidazolidin-1-yl)-pent-4-enoic acid,
- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-{3-[6-(4-methoxy-benzylamino)-pyridin-2-  
yl]-propyl}-2-oxo-imidazolidin-1-yl)-propionic acid,

- 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-{3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl}-2-oxo-imidazolidin-1-yl)-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-(3-{3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl}-2-oxo-imidazolidin-1-yl)-propionic acid,
- 5 3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 10 3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3(R)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 15 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 20 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 25 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 30 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 35

- 3-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15 3-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3 3-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 3(R)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 15 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 20 3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 25 3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 35 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

5 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

10

3-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

15 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

20 3-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

25

3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

30 3-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

20. The compound of Claim 19 selected from the group consisting of

- 5 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3(S)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(S)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(R)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3(R)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 bis(trifluoroacetate) salt,
- 3(R)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(S)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35



3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 5 3(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

10

3(S)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 15 3(R)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 20 3(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25

3(R)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 30 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3(S)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 5 3(S)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 10 3(R)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 15 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 20 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(R)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 35

- 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 25 3(R)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 35

- 3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
15 [1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 25 3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
30 [1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 35 and the pharmaceutically acceptable salts thereof.

21. The compound of Claim 20 selected from the group consisting of

- 5 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(S)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid, bis(trifluoroacetate) salt,
- 3(S)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(S)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,
- 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,
- 35

3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-  
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-  
5 2-yl)propyl]imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

22. A pharmaceutical composition comprising a  
10 compound according to Claim 1 and a pharmaceutically acceptable  
carrier.

23. A pharmaceutical composition made by combining a  
compound according to Claim 1 and a pharmaceutically acceptable  
15 carrier.

24. A process for making a pharmaceutical composition  
comprising combining a compound according to Claim 1 and a  
pharmaceutically acceptable carrier.

20

25. The composition of Claim 22 which further comprises  
an active ingredient selected from the group consisting of

- a) an organic bisphosphonate or a pharmaceutically  
acceptable salt or ester thereof,
- 25 b) an estrogen receptor modulator,
- c) a cytotoxic/antiproliferative agent,
- d) a matrix metalloproteinase inhibitor,
- e) an inhibitor of epidermal-derived, fibroblast-derived, or  
platelet-derived growth factors,
- 30 f) an inhibitor of VEGF,
- g) an inhibitor of Flk-1/KDR, Flt-1, Tck/Tie-2, or Tie-1,
- h) a cathepsin K inhibitor, and
- i) a prenylation inhibitor, such as a farnesyl transferase  
inhibitor or a geranylgeranyl transferase inhibitor or a dual  
35 farnesyl/geranylgeranyl transferase inhibitor;

and mixtures thereof.

26. The composition of Claim 25 wherein said active ingredient is selected from the group consisting of

- 5           a) an organic bisphosphonate or a pharmaceutically acceptable salt or ester thereof,  
          b) an estrogen receptor modulator, and  
          c) a cathepsin K inhibitor;  
          and mixtures thereof.

10

27. The composition of Claim 26 wherein said organic bisphosphonate or pharmaceutically acceptable salt or ester thereof is alendronate monosodium trihydrate.

15

28. The composition of Claim 25 wherein said active ingredient is selected from the group consisting of

- a) a cytotoxic/antiproliferative agent,  
          b) a matrix metalloproteinase inhibitor,  
          c) an inhibitor of epidermal-derived, fibroblast-derived, or  
20           platelet-derived growth factors,  
          d) an inhibitor of VEGF, and  
          e) an inhibitor of Flk-1/KDR, Flt-1, Tck/Tie-2, or Tie-1;  
          and mixtures thereof.

25

29. A method of eliciting an integrin receptor antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound according to Claim 1.

30

30. The method of Claim 29 wherein the integrin receptor antagonizing effect is an  $\alpha v\beta 3$  antagonizing effect.

31. The method of Claim 30 wherein the  $\alpha v\beta 3$  antagonizing effect is selected from the group consisting of inhibition of

bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, and tumor growth.

32. The method of Claim 31 wherein the  $\alpha v \beta 3$   
5 antagonizing effect is the inhibition of bone resorption.

33. The method of Claim 29 wherein the integrin receptor antagonizing effect is an  $\alpha v \beta 5$  antagonizing effect.

10 34. The method of Claim 33 wherein the  $\alpha v \beta 5$  antagonizing effect is selected from the group consisting of inhibition of restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, and tumor growth.

15 35. The method of Claim 29 wherein the integrin receptor antagonizing effect is a dual  $\alpha v \beta 3 / \alpha v \beta 5$  antagonizing effect.

20 36. The method of Claim 35 wherein the dual  $\alpha v \beta 3 / \alpha v \beta 5$  antagonizing effect is selected from the group consisting of inhibition of bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, and tumor growth.

25 37. The method of Claim 29 wherein the integrin antagonizing effect is an  $\alpha v \beta 6$  antagonizing effect.

38. The method of Claim 37 wherein the  $\alpha v \beta 6$  antagonizing effect is selected from the group consisting of angiogenesis, inflammatory response, and wound healing.

30 39. A method of eliciting an integrin receptor antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 22.



40. A method of treating or preventing a condition mediated by antagonism of an integrin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 22.

5

41. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 22.

10

42. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 26.

15

43. A method of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 28.

20

44. A method of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound according to Claim 1 in combination with radiation therapy.

# INTERNATIONAL SEARCH REPORT

Internati	Application No
PCT/US 98/26568	

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 C07D401/06 C07D401/14 C07D519/00 A61K31/415  
A61K31/44 //(C07D519/00,498:00,471:00),(C07D519/00,491:00,  
471:00),(C07D519/00,471:00,471:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 051 829 A (A. NATTERMANN & CIE. GMBH) 19 May 1982 see claims 1,20; examples 1,18,48	1,22
A	WO 95 17397 A (MERCK & CO., INC.) 29 June 1995 cited in the application see claims 1,4,5	1,22
A	WO 95 32710 A (MERCK & CO., INC.) 7 December 1995 cited in the application see claims 11-14,17	1,22
A	WO 91 18888 A (G. D. SEARLE & CO.) 12 December 1991 see claims 1,22; examples 7,9,10,24,27	1,22
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

7 April 1999

Date of mailing of the international search report

13/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Hass, C

# INTERNATIONAL SEARCH REPORT

Internat. Application No.  
PCT/US 98/26568

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 93 16674 A (SMITHKLINE BEECHAM CORP.) 2 September 1993 see claims 26-28	1,22
A	US 4 460 595 A (D. R. ADAMS ET AL.) 17 July 1984 see claim 1; tables 3,6	1,22
A	EP 0 006 718 A (BEECHAM GROUP PLC) 9 January 1980 see claims 1-3,11	1,22

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 26568

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 29-44  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Although claims 29-44 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

---

Claims Nos.: 29-44

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No

PCT/US 98/26568

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 51829	A	19-05-1982	DE 3042466 A AT 10626 T DK 495781 A GR 74703 A IE 52079 B JP 57108073 A US 4349684 A ZA 8107696 A	16-06-1982 15-12-1984 12-05-1982 05-07-1984 10-06-1987 05-07-1982 14-09-1982 27-10-1982
WO 9517397	A	29-06-1995	US 5416099 A AU 1441695 A	16-05-1995 10-07-1995
WO 9532710	A	07-12-1995	AU 701776 B AU 2586895 A CA 2190870 A EP 0760658 A JP 10501222 T US 5741796 A	04-02-1999 21-12-1995 07-12-1995 12-03-1997 03-02-1998 21-04-1998
WO 9118888	A	12-12-1991	AU 7903091 A CA 2083735 A EP 0531382 A	31-12-1991 26-11-1991 17-03-1993
WO 9316674	A	02-09-1993	AU 3664593 A AU 701541 B AU 4186797 A AU 703236 B AU 6482698 A CA 2129897 A EP 0626969 A US 5648373 A US 5663053 A CA 2061061 A	13-09-1993 28-01-1999 08-01-1998 18-03-1999 16-07-1998 12-08-1993 07-12-1994 15-07-1997 02-09-1997 12-08-1993
US 4460595	A	17-07-1984	GB 1598073 A AU 509364 B AU 3161277 A CA 1097660 A CA 1101870 A CH 638190 A DE 2755727 A DK 551677 A FI 773813 A FR 2374313 A IE 46118 B JP 53082781 A NL 7713973 A SE 7714372 A US 4367338 A BE 861957 A ZA 7707336 A	16-09-1981 08-05-1980 21-06-1979 17-03-1981 26-05-1981 15-09-1983 29-06-1978 19-06-1978 19-06-1978 13-07-1978 23-02-1983 21-07-1978 20-06-1978 19-06-1978 04-01-1983 16-06-1981 25-10-1978
EP 6718	A	09-01-1980	AU 4806879 A JP 55004385 A	20-12-1979 12-01-1980